

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07K 1/00, 7/06, A61K 37/02</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/22335</b> <b>(43) International Publication Date:</b> 11 November 1993 (11.11.93)
<b>(21) International Application Number:</b> PCT/US93/03933 <b>(22) International Filing Date:</b> 27 April 1993 (27.04.93) <b>(30) Priority data:</b> 07/876,625 30 April 1992 (30.04.92) US <b>(71) Applicant:</b> COR THERAPEUTICS, INC. [US/US]; 256 East Grand Avenue, Suite 80, South San Francisco, CA 94080 (US). <b>(72) Inventors:</b> SWIFT, Robert, L. ; 806 Prowshead Lane, Foster City, CA 94404 (US). DU MEE, Charles, P. ; 18 Coral Lane, Foster City, 94404 (US). RANDOLPH, Anne, E. ; 1337 Drake Avenue, Burlingame, CA 94010 (US).		<b>(74) Agents:</b> PARK, Freddie, K. et al.; Morrison & Foerster, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US). <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> STABLE POLYPEPTIDE COMPOSITION  <b>(57) Abstract</b>  A (injectable biologically active) polypeptide is stabilized by dissolving said polypeptide forming a liquid solution in citrate buffer of about pH 5.0-5.5.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

STABLE POLYPEPTIDE COMPOSITIONField of the Invention

The present invention relates to stable polypeptide compositions and methods of preparing such stable  
10 compositions.

Background of the Invention

Polypeptides possess chemical and physical properties that present special problems which can cause  
15 instability in storage and delivery, particularly of biologically active polypeptides. Usually polypeptides are lyophilized to form solid compositions. Often, additives are introduced into polypeptide formulations in an effort to increase stability. Such additives have  
20 included salts of ionic compounds, polyalcohols, but this is less effective at higher concentrations of such additive, and detergents, both nonionic and anionic, particularly for transdermal and intranasal administration. Many of the instability problems of  
25 protein pharmaceuticals are discussed in Manning et al., Pharm. Res. (1989) 6(11):903-918.

European patents 311,950, 286,830 and 288,891 and U.S. patents 4,361,510 and 4,470,968 describe separation processes that use of one or usually more than one of  
30 various additives such as 3-propiolactone; various Tris, hepes, glycine or citrate buffers, polyols or alkanols; salts, such as sodium chloride or sodium, potassium or lithium citrate, in preparing or purifying proteins but with a pH above at least 5.5 and preferably usually 6, if  
35

-2-

not 7-8. Usually the protein is removed from these additives and preferably lyophilized.

However, many polypeptides are useful as biologically active materials which are most useful when injected directly into an animal. This is the case for polypeptides useful as therapeutic agents for the treatment and prevention of platelet-associated ischemic disorders initiated by atherosclerosis and arteriosclerosis.

In treatment of some life-threatening results of these disorders it can be necessary to inject the patient with a drug concentrate as quickly as possible (bolus treatment) and/or followed by prolonged (infusion) treatment with a lower drug concentration. Accordingly, the drug, which may be a polypeptide, must be stored conveniently for immediate use by hospital personnel and paramedics. This means it must be storage-stable for long periods, even at room temperature, and already in an injectable liquid formulation, not a solid which must be dissolved.

Because the drug is injectable, it must be formulated as simply as possible to be as compatible as possible with the patient so as to not induce other traumas or interfere with other drugs being given to the patient. The ideal formulation would avoid other additives normally required for stabilization and be a ready-to-use liquid concentrate.

#### Summary of the Invention

The present invention is directed to a method of stabilizing substantially pure polypeptide comprising forming a liquid solution consisting essentially of a substantially pure polypeptide in a citrate buffer, said solution having a pH of from about 5.0 to about 5.5, to stabilize said polypeptide. Preferably, the polypeptide

-3-

is an injectable, biologically active polypeptide so that the stabilized composition can be used as a resulting therapeutically effective solution.

The invention also includes the resulting useful therapeutic compositions of an injectable biologically active, substantially pure polypeptide consisting essentially of a biologically effective amount of a substantially pure polypeptide in a liquid solution of a citrate buffer, said solution having a pH of from about 5.0 to about 5.5. Such compositions are capable of remaining very stable, for example, at about 4 °C, for 7, 34 or 49 days prior to injection, are still stable at up to about 50°C and have greater stability at about 70°C than compositions having a pH above 5.5. The compositions of the invention also remain stable (and injectable) at about -15° C to about 30° C for at least 90 days, preferably for at least 18 months and have improved stability at higher temperatures.

The resulting liquid composition can be aseptically introduced into sterile delivery vial, bag or bottle and sealed ready for injection on attachment of an injection device, such as a hypodermic needle or intravenous tube.

#### Detailed Description of the Invention

The invention provides a method of stabilizing a substantially pure polypeptide in a liquid solution to be stable on storage at about -15° C to about 30°C for at least 90 days and, preferably, at least 18 months or to have improved stability at higher temperatures. The polypeptide is dissolved in the citrate buffer to form the solution.

By "substantially pure" is meant a polypeptide that has been purified to a pharmaceutically acceptable level of purity and preferably is essentially purified to homogeneity.

**SUBSTITUTE SHEET**

-4-

Polypeptides soluble in aqueous media are well known to those of skill in the art, including as described in U.S. Patent 4,532,212 and includes derivatives such as enzymes and the like. Pharmaceutically useful  
5 polypeptides are well known to those of skill in the art as discussed in manning, et al. Pharmaceutical Research, 6 (11): 903-918 (1989).

By "biologically active" is meant any polypeptide having an effect on a biological system. Such  
10 polypeptides include pharmaceutically active polypeptides, such as interferon, insecticidally active polypeptides, such as insecticidal Bacillus thuringiensis endotoxins, and the like. A wide variety of polypeptides are known in the art which have an effect on a biological  
15 system, in the case of pharmaceuticals the effect is therapeutic. In other cases the effect can be insecticidal, fungicidal or the like.

By "injectable" is meant that the polypeptide can be injected into a patient and a wide variety of such  
20 polypeptides are known to those of skill in the art.

By "biologically compatible citrate buffer" is meant a citrate buffer prepared from ingredients that do not themselves have an undesirable or adverse effect on a biological system. Such buffer components are well known  
25 to those of skill in the pharmaceutical art.

By "biological system" is meant a living entity including plants, animals or a living part thereof, such as an organ or cell. The preferred biological system is a mammalian system, especially a human system.

30 By "stability" or "improved stability" is meant herein that in the liquid solution of the invention, a high amount (e.g., by weight) of the substantially pure polypeptide remains substantially intact or more intact (that is physically and chemically stable and therefore  
35 biological activity when initially present is also

SUBSTITUTE SHEET

-5-

stabilized) after a period of time under conditions of exposure to temperatures below 50°C, preferably below 30°C and has improved stability even at about 70°C, and to light, oxygen and the storage container used for injectable biologically active substantially pure polypeptides. This stability can be evaluated by conventional assay methods applicable to purity, weight or size of polypeptides. These include not only visual evaluations, such as discoloration, transparency and precipitation, but can include assays normally applied to separate polypeptides from each other and from other materials, such as chromatography, e.g., reversed phase high performance liquid chromatography (HPLC), and ultraviolet (UV) analysis. Conventional biological assays can be used when the polypeptide is a biologically active one.

By "citrate buffer" is meant a conventional citrate solution to which large amounts of strong acid or base can be added with only very small resultant change in the hydrogen-ion concentration. The citrate buffer is prepared by methods known in the art by adding a pH-neutral strong electrolyte or a strong base to a citric acid solution. Suitable electrolytes include sodium chloride, sodium citrate, potassium nitrate, sodium hydrogen sulfate and the like. Suitable strong bases include sodium hydroxide, calcium hydroxide and the like, sodium hydroxide is preferred. Adjustment in ionic strength, pH concentration of polypeptide are made in the normal manner for making pH-adjusted isotonic solutions. Additions of ingredients to prepare the citrate buffer and compositions of the invention can be made in any order but preferably, the polypeptide is added to injectable grade water and then the buffer ingredients are added or especially when the polypeptide is added to a citric acid solution. In one embodiment, a solution of

SUBSTITUTE SHEET

-6-

a concentration of about 200 mg of polypeptide per ml in 1.0 M citric acid is formed, diluted to 85% of the final volume with water, the pH is adjusted to 5.0 to 5.5 using sodium hydroxide and then diluted to the final volume and concentration with water.

Additional ingredients which are conventionally employed in (pharmaceutical) preparations, such as glycine and other salts, can be present but are not essential ingredients in the compositions of the invention.

While a wide variety of biologically compatible buffers are known in the art, they are usually used with other additives at a pH above 5.5 and usually at a pH above 7, and even then lyophilized to a dry powder. However, it has now been found that citrate buffer solutions of a (biologically active) polypeptide having a pH of from about 5.0 to about 5.5 are unexpectedly very storage-stable over long periods of time without the addition of other ingredients or without other treatments such as lyophilization. In a preferred embodiment, the pH of the solution is from about 5.25.

The concentration of the substantially pure polypeptide in the citrate buffer is a stabilizing effective amount and can vary depending on a variety of factors including the specific polypeptide, pH and the specific buffer. The stabilizing effective amount of the citrate buffer and the concentration of polypeptide in a stabilized composition of the invention can readily be determined by those of skill in the protein formulation art. As a general matter, the concentration of the substantially pure polypeptide can be from about 0.01 to about 200 mg/ml and is usually about 0.01 to about 10 mg/ml. For injectable and/or biologically active polypeptides, the concentration in the composition can further be adjusted as can readily be determined by those

**SUBSTITUTE SHEET**



-7-

of skill in the art to correspond to an effective dosage for the intended biological effect. For example, when the polypeptide is Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate or a cyclic form thereof, the  
5 concentration usually corresponds to a dosage of from about 0.5 to about 5 mg/ml, preferably from about 0.5 to about 2.0 mg/ml. A 0.5 mg/ml dosage form, which is a direct infusion pre-mix is a bolus followed by an infusion. The bolus is about 10 to about 500, preferably  
10 about 30-300 µg/Kg (about 1-40 mg, preferably from about 2-20 mg for a 70Kg patient) and the infusion is 0.02-2 µg/Kg/min, but probably 0.5-1.0 µg/Kg/min (50-100 mg/day for a 70Kg patient). Administration can be by injection from a bolus up to 7 days, preferably for about 12 hours  
15 to about 3 days depending on the indication.

Preferably, the polypeptide is any biologically active, substantially pure polypeptide that can be injected into a patient for treatment can be stabilized by the method of the invention. The substantially pure  
20 polypeptides include linear and cyclic polypeptides of at least two amino acid residues and therefore includes simple linear dipeptides and large protein molecules. Preferably, the polypeptide is a synthetic or recombinant polypeptide.

25 In one preferred embodiment of the invention, the polypeptide contains up to 10 amino acid residues. An especially preferred polypeptide is Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate or a cyclic form thereof.


30 The substantially pure polypeptide can contain the usual polypeptide modifications such as disulfide bonds, ionic bonds, glycosylation and the like.

35

SUBSTITUTE SHEET


-8-

In one embodiment of the invention, the polypeptide has the formula



5

wherein  $K^*$  has the formula



10

wherein (G/Sar) is G or Sar;

wherein each  $R^1$  is independently H, alkyl(1-6C), or at most one  $R^1$  is  $R^2-C=NR^3$ ,


wherein  $R^2$  is H, alkyl(1-6C) or is a substituted or unsubstituted phenyl or benzyl residue, or is  $NR^4_2$  in

15

which each  $R^4$  is independently H or alkyl(1-6C), and

$R^3$  is H, alkyl(1-6C), phenyl or benzyl, or  $R^2-C=NR^3$  is a radical selected from the group consisting of

20



25

where  $m$  is an integer of 2-3, and each  $R^5$  is independently H or alkyl(1-6C);

and wherein one or two  $(CH_2)$  may be replaced by O or S provided said O or S is not adjacent to another heteroatom;

30

$AA_1$  is a small, neutral (polar or nonpolar) amino acid and  $n1$  is an integer of 0-3;

$AA_2$  is a neutral, nonpolar large (aromatic or nonaromatic) or a polar aromatic amino acid and  $n2$  is an integer of 0-3;

35

$AA_3$  is a proline residue or a modified proline residue and  $n3$  is an integer of 0-1;

SUBSTITUTE SHEET

-9-

AA<sub>4</sub> is a neutral, small amino acid or the N-alkylated form thereof and n<sub>4</sub> is an integer of 0-3;

each of X<sub>1</sub> and X<sub>2</sub> is independently a residue capable of forming a bond between X<sub>1</sub> and X<sub>2</sub> to obtain a cyclic compound as shown; and

each of Y<sub>1</sub> and Y<sub>2</sub> is independently a noninterfering substituent or may be absent;

wherein one or more peptide linkages may optionally be replaced by a linkage selected from the group consisting of -CH<sub>2</sub>NH-, -CH<sub>2</sub>S-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- (cis and trans), -COCH<sub>2</sub>-, -CH(OH)CH<sub>2</sub>- and -CH<sub>2</sub>SO-.

Suitable peptides of the invention include those wherein Y<sub>1</sub> is H, acyl, or a peptide residue or derivatized form thereof or is absent and Y<sub>2</sub> is OH, NH<sub>2</sub> or a peptide residue or derivatized form thereof or is absent, Y<sub>2</sub> is NH<sub>2</sub>-A-NH<sub>2</sub> or is absent, X<sub>1</sub> and X<sub>2</sub> are selected from the group consisting of cysteine (C), mercaptopropionyl (Mpr) and penicillamine (Pen), AA<sub>1</sub> is G and n<sub>1</sub> is 0 or 1, AA<sub>2</sub> is selected from the group consisting of W, F, L, Y, and V and K\* is K, Har, acetimidyl-Lys or phenylimidyl-Lys. In one embodiment, the polypeptides have the above formula with the proviso that if n<sub>3</sub> is 0; either:

- 1) the sum of n<sub>2</sub> and n<sub>4</sub> must be at least 2; or
- 2) K\* must be other than Har or K; or
- 3) at least one of X<sub>1</sub> and X<sub>2</sub> must be other than cys (C), penicillamine (Pen), or 2-amino-3,3-cyclopentanemethylene-3-mercaptopropionic acid (APmp); or
- 4) Y<sub>1</sub> or Y<sub>2</sub> must comprise at least one amino acid residue; or
- 5) one or more linkages is replaced by said alternate linkage.

Examples of suitable peptides include  
 Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>  
 Mvl-K-G-D-W-P-C-NH<sub>2</sub>

SUBSTITUTE SHEET

-10-


- Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 5 Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub>  
 10 Mpr-K-G-D-W-P-C-NH<sub>2</sub>  
 or cyclic forms thereof.

Such peptides are described in copending USSN  
 07/483,229, filed 2/20/90 and PCT/US90/03417 published as  
 W090/15620, each incorporated herein by reference and  
 15 described in further detail in the following 35 pages and  
 examples 4-14.

"Alkyl" is conventionally defined as a straight or  
 branched chain or cyclic hydrocarbyl residue of the  
 indicated number of carbon atoms such as methyl, ethyl,  
 20 isopropyl, N-hexyl, 2-methylbutyl, cyclohexyl and the  
 like.

The benzyl and phenyl residues represented by R<sup>2</sup> may  
 be unsubstituted, or may be substituted by noninterfering  
 substituents. Preferred substitution patterns are those  
 25 wherein only one substituent is bound to the aromatic  
 nucleus, preferably in the 4-position. Preferred  
 substituents are electron donating substituents such as  
 alkyl (1-6C), especially ethyl or methyl, or phenyl.

Preferred embodiments of K<sup>\*</sup> include the residues of  
 30 lysine, homoarginine, formylhomoarginine, ornithine,  
 acetimidyl lysine, N<sup>G</sup>N<sup>G</sup> ethylene-homoarginine, and  
 phenylimidyl lysine. The phenylimidyl lysyl residue, for  
 example, has the formula:



35

-11-

As the essential feature of the preferential inhibition of binding appears to reside in the substitution of K\* for R of RGD<sub>X</sub>, one class of peptides or peptide-related compounds for use in the compositions of the invention comprises naturally occurring platelet aggregation inhibitors (PAI) which ordinarily contain RGD<sub>X</sub> in the binding sequence whereby these forms are modified by substituting K\* for R in this sequence. Included are the native peptides having this substitution, as well as their fragments of sufficient length to be effective in selectively inhibiting the binding of adhesive proteins to GP IIb-IIIa and fragments or full-length peptides which have irrelevant substitutions in positions of the peptide which do not destroy this activity. For the most part, the fragments will contain residues corresponding to the length of a peptide chain of at least 7 amino acids if the conformation is controlled by, for example, cyclization, and are of greater length if there is no such conformational control. In general, aside from the K\*GD<sub>X</sub> required sequence, there may be 1-10, preferably 14, and preferably 1-3 amino acid substitutions in the non-K\*GD<sub>X</sub> portion of the peptides.

Additionally, the G of RGD<sub>X</sub> or K\*GD<sub>X</sub> can be replaced by a sarcosine residue.

In addition, one or more of the peptide bonds can be optionally replaced by substitute linkages such as those obtained by reduction or elimination. Thus, one or more of the -CONH- peptide linkages can be replaced with other types of linkages such as -CH<sub>2</sub>NH-, -CH<sub>2</sub>S-, CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- (cis and trans), -COCH<sub>2</sub>-, -CH(OH)CH<sub>2</sub> and -CH<sub>2</sub>SO-, by methods known in the art. The following references describe preparation of peptide analogs which include these alternative-linking moieties: Spatola, A.F., Vega Data (March 1983), Vol. 1, Issue 3, "Peptide Backbone

SUBSTITUTE SHEET

-12-

- Modifications" (general review); Spatola, A.F. in "Chemistry and Biochemistry of Amino Acids, Peptides and Proteins," B. Weinstein, eds., Marcel Dekker, New York, p. 267 (1983) (general review); Morley, J.S., Trends Pharm Sci (1980) pp. 463-468 (general review); Hudson, D. et al. Int J Pept Prot Res (1979) 14:177-185 (-CH<sub>2</sub>NH-, CH<sub>2</sub>CH<sub>2</sub>-); Spatola, A.F. et al., Life Sci (1986) 38:1243-1249 (-CH<sub>2</sub>-S); Hann, M.M. J Chem Soc Perkin Trans I (1982) 307-314 (-CH-CH-, cis and trans); Almquist, R.G., et al., J Med Chem (1980) 23: 1392-1398 (-COCH<sub>2</sub>); Jennings-White, C. et al. Tetrahedron Lett (1982) 23:2533 (-COCH<sub>2</sub>-); Szelke, M., et al., European Appln. EP 45665 (1982) CA: 97:39405 (1982) (-CH(OH)CH<sub>2</sub>-); Holladay, M.W. et al. Tetrahedron Lett (1983) 24:4401-4404 (-C(OH)CH<sub>2</sub>-); and Hruby, V.J. Life Sci (1982) 31:189-199 (-CH<sub>2</sub>-S-). Particularly preferred is -CH<sub>2</sub>NH-.

Examples of fragments and/or modified forms of the naturally-occurring snake venom PAI include

[E<sup>28</sup>, L<sup>41</sup>, C<sup>64</sup>]barbourin(28-73) of the sequence

1 ECADGLCCDQCRFLKKGTVCRAKGDWNDDTCTGQSCDCPRNGLYG 46  
28 73

and [K<sup>29</sup>]eristicophin(4-51) of the sequence

4 EEPCATGPCCRRCKFKRAGKVCRAKGDWNNDYCTGKSCDCPRNPWNG. 51  
4 51

In this notation, the size of the fragment is noted in parentheses after the name by the numbers of the amino acids which are included in the fragment, and the bracketed prefix letters and numbers indicate amino acid substitutions at the numbered positions in the native full-length peptide. Thus, for the barbourin fragment above, the length of the fragment spans residues 28-73 inclusive of the native sequence and the amino acids originally in positions 28, 41 and 64 of the numbered native sequence

-13-

have been replaced by Glu (E), Leu (L), and Cys (C), respectively.

As additional examples, the arginine of the RGD sequence appearing in trigramin, elegantin, albolabrin, crotatroxin, flavoviridin, echistatin, bitistatin, viridin, molossin, lutosin, basilicin, applagin, halysin, horridin, tergeminin, lachesin, cotiarin, cereberin, jararacin, kistrin, eristicophin, bitan-a, and ruberin/oreganin can be replaced by a K<sup>\*</sup> residue to provide specifically active PAIs with a preferential affinity for GP IIb-IIIa. In addition, shortened forms of these peptides, containing at least 20, preferably at least 30, and more preferably at least 40, amino acids, can be prepared from the native peptide or in this modified form. In addition, or in the alternative, 1-10, preferably 1-4, amino acids irrelevant to the RGD/K<sup>\*</sup>GD sequence can be substituted or modified, preferably with conservative amino acid substitutions. By conservative amino acid substitutions is meant, for example, substitution of an acidic amino acid residue for an acidic amino acid residue, neutral for neutral, basic for basic, etc., as is further described hereinbelow.

Still an additional group of examples includes that wherein the glycyl residue of RGD or K<sup>\*</sup>GD can be replaced by a sarcosyl residue with retention of activity. Thus, the active PAIs which are isolated and/or modified in other ways as described above may further be modified by this substitution.

While fragments and/or modified PAIs from snake venom can be included among the Fg/vWF/GP IIb-IIIa binding-specific compounds by replacing RGD by K<sup>\*</sup>GD, in additional embodiments specifically active peptides are based on compatible extensions of the K<sup>\*</sup>GD sequence per se.

Y<sub>1</sub> and Y<sub>2</sub> can be peptide extensions of 0-25 amino acid residues and can be in derivatized form. The Y<sub>1</sub> N-terminal extensions can, for example, be acetylated or otherwise

SUBSTITUTE SHEET

-14-

acylated; the  $Y_2$  C-terminal extension can be amidated with  $NH_2$  or with a primary or secondary amine of the formula  $R-NH_2$  or  $R_2NH$  wherein each  $R$  is independently a lower alkyl of 1-4C, such as methyl, n-butyl, or t-butyl.  $Y_1$  can also  
5 be (H) or acyl (1-4C);  $Y_2$  can be (OH),  $NH_2$  or an amine as above. Where the compound of formula (1) is a simply cyclic peptide,  $Y_1$  and  $Y_2$  are absent.

$X_1$  and  $X_2$  are typically amino acid residues capable of cyclization such as, for example and most preferably,  
10 cysteine residues capable of forming a disulfide ring. However, other residues capable of forming disulfide or other linkages can also be used -- for example, the Pen (penicillamine) residue described by Pierschbacher et al. (supra) or the Mpr (mercapto propionyl) or Mvl  
15 (mercaptovaleryl) residue. Other types of covalent linkages for cyclization envisioned include peptide linkages, as for example, an amide formed between the side-chain amino group of a lysyl residue with a side-chain carboxyl group of a glutamyl residue and ester linkages,  
20 such can be formed between a side-chain alcohol of a threonine residue with a side-chain carboxyl of an aspartyl residue. Any compatible residue capable of forming peptide bonds with the remainder of the chain (or modified peptide bonds as described above) and capable of covalent bond  
25 formation to effect cyclization can be used. This includes, for example, simply cyclic peptides, wherein a peptide bond is directly formed between the  $NH^2$  at the N-terminus and the COOH at the C-terminus.

As described above, one or more of the indicated  
30 peptide bonds may be replaced by a substitute linkage such as  $-CH_2NH-$ ,  $-CH_2S-$ ,  $CH_2CH_2-$ ,  $-CH=CH-$  (cis and trans),  $-COCH_2-$ ,  $-CH(OH)CH_2-$  and  $-CH_2SO-$ .

In the designation of the amino acid residues  $AA_1-AA_4$  above, description has been made on the basis of a  
35 classification method, wherein amino acid residues can be

SUBSTITUTE SHEET



-15-

generally subclassified into four major subclasses. This classification is also shown diagrammatically hereinbelow.

5        Acidic: The residue has a negative charge due to loss of H ion at physiological pH and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH.

10        Basic: The residue has a positive charge due to association with H ion at physiological pH and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH.

15        Neutral/nonpolar: The residues are not charged at physiological pH and the residue is repelled by aqueous solution so as to seek the inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. These residues are also  
20        designated "hydrophobic" herein.

      Neutral/polar: The residues are not charged at physiological pH, but the residue is attracted by aqueous solution so as to seek the outer positions in the conformation of a peptide in which it is contained when the  
25        peptide is in aqueous medium.

      It is understood, of course, that in a statistical collection of individual residue molecules some molecules will be charged, and some not, and there will be an attraction for or repulsion from an aqueous medium to a  
30        greater or lesser extent. To fit the definition of "charged", a significant percentage (at least approximately 25%) of the individual molecules are charged at physiological pH. The degree of attraction or repulsion required for classification as polar or nonpolar is  
35        arbitrary, and, therefore, amino acids specifically

SUBSTITUTE SHEET

-16-

contemplated have been specifically classified as one or the other. Most amino acids not specifically named can be classified on the basis of known behavior.

Amino acid residues can be further subclassified as cyclic or noncyclic, and aromatic or nonaromatic, self-explanatory classifications with respect to the side chain substituent groups of the residues, and as small or large. The residue is considered small if it contains a total of 4 carbon atoms or less, inclusive of the carboxyl carbon. Small residues are, of course, always nonaromatic.

For the naturally occurring protein amino acids, subclassification according to the foregoing scheme is as follows (see also the diagram below).


- 15      Acidic: Aspartic acid and Glutamic acid;  
         Basic/noncyclic: Arginine, Lysine;  
         Basic/cyclic: Histidine;  
         Neutral/polar/small: Glycine, Serine and Cysteine;  
20      Neutral/polar/large/nonaromatic: Threonine,  
         Asparagine, Glutamine;  
         Neutral/polar/large/aromatic: Tyrosine;  
         Neutral/nonpolar/small: Alanine;  
25      Neutral/nonpolar/large/nonaromatic: Valine,  
         Isoleucine, Leucine, Methionine;  
         Neutral/nonpolar/large/aromatic: Phenylalanine, and  
         Tryptophan.

30      The gene-encoded amino acid proline, although technically within the group neutral/nonpolar/large/cyclic and nonaromatic, is a special case due to its known effects on the secondary conformation of peptide chains, and is not, therefore, included in this defined group, but is classified separately. AA<sub>3</sub> is designated a  
35      proline residue or a "modified proline residue."

SUBSTITUTE SHEET

-17-

Proline, as is understood, is a five-membered nitrogen heterocycle with a carboxyl group in the 2-position. Modified proline residues are all nitrogen five or six-membered heterocycles with carboxyl groups in the position alpha to the nitrogen; additional heterocyclic atoms may also be included in the ring. Thus, modified proline residues include residues of pipecolic acid (2-carboxypiperidine, abbreviated Pip) and thiazolidine (Thz). Thus, proline or modified proline residues are of the formula



wherein one or two of the methylene groups may be replaced by NR, S, or O and where any ring nitrogen can optionally be substituted with R as a noninterfering substituent such as alkyl (1-6C).

Certain commonly encountered amino acids, which are not encoded by the genetic code, include, for example, beta-alanine (beta-ala), or other omega-amino acids, such as 3-amino propionic, 4-amino butyric and so forth, alpha-aminoisobutyric acid (Aib), sarcosine (Sar), ornithine (Orn), citrulline (Cit), homoarginine (Har), t-butylalanine (t-BuA), t-butylglycine (t-BuG), N-methylisoleucine (N-MeIle), phenylglycine (Phg), and cyclohexylalanine (Cha), norleucine (Nle), cysteic acid (Cya); pipecolic acid (Pip), thiazolidine (Thz), 2-naphthyl alanine (2-Nal) and methionine sulfoxide (MSO). These also fall conveniently into particular categories.

Based on the above definition,  
 Sar and beta-ala are neutral/nonpolar/small;  
 t-Bua, t-BuG, N-MeIle, Nle and Cha are neutral/  
 nonpolar/large/nonaromatic;  
 Cya is acidic;

SUBSTITUTE SHEET

-18-

Cit, Acetyl Lys, and MSO are  
neutral/polar/large/nonaromatic;

2-Nal and Phg are  
neutral/nonpolar/large/aromatic; and

5           Pip and Thz are modified proline residues.

The foregoing may be shown diagrammatically as  
follows:

10

15

20

25

30

35

**SUBSTITUTE SHEET**



-20-

The various omega-amino acids are classified according to size as neutral/nonpolar/small (beta-ala, i.e., 3-aminopropionic, 4-aminobutyric) or large (all others).

5 Other amino acid substitutions for those encoded in the gene can also be included in peptide compounds within the scope of the invention and can be classified within this general scheme.

10 In the formulas representing selected specific embodiments of the present invention, the amino- and carboxyl-terminal groups, although often not specifically shown, will be understood to be in the form they would assume at physiological pH values, unless otherwise specified. Thus, the N-terminal  $H^+$  and C-terminal  $O^-$  at  
15 physiological pH are understood to be present though not necessarily specified and shown, either in specific examples or in generic formulas. Of course, the basic and acid addition salts including those which are formed at nonphysiological pH values are also included. Unless  
20 otherwise noted, the residues are in the L-form; in generic formulas, the specified residues can be either L- or D-. Generally, the polypeptides of the invention have 0, 1, or 2 D-residues included, preferably 0 or 1, most preferably 0. In the polypeptides shown, each encoded  
25 residue where appropriate is represented by a single letter designation, corresponding to the trivial name of the amino acid, in accordance with the following conventional list:

30	<u>Amino Acid</u>	<u>One-Letter Symbol</u>
	Alanine	A
	Arginine	R
	Asparagine	N
	Aspartic acid	D
	Cysteine	C
	Glutamine	Q
35	Glutamic acid	E

-21-

	Glycine	G
	Histidine	H
	Isoleucine	I
	Leucine	L
	Lysine	K
	Methionine	M
5	Phenylalanine	F
	Proline	P
	Serine	S
	Threonine	T
	Tryptophan	W
	Tyrosine	Y
	Valine	V
10	Pyroglutamic acid	Z

The amino acids not encoded genetically are abbreviated as indicated above.

15 In the specific peptides shown in the present application, the L-form of any amino acid residue having an optical isomer is intended unless otherwise expressly indicated by a dagger superscript (<sup>†</sup>). While the residues of the peptides are normally in the natural L optical isomer form, one or two, preferably one, amino  
20 acid may be replaced with the optical isomer D form.

Free functional groups, including those at the carboxy- or amino-terminus, can also be modified by amidation, acylation or other substitution, which can, for example, change the solubility of the compounds  
25 without affecting their activity.

In forming amidated peptides of the present invention, the analog compounds can be synthesized directly, for example using BOC-AA<sub>x</sub>-pMBHA-Resin or Boc-AA<sub>x</sub>-BHA-Resin, wherein AA<sub>x</sub> is the selected carboxy-  
30 terminal amino acid of the desired peptide as described in further detail below. Alternatively, the peptides can be chemically or enzymatically amidated subsequent to peptide synthesis using means well known to the art, or prepared by standard solution-phase peptide synthesis  
35 protocols.

**SUBSTITUTE SHEET**

-22-

Certain embodiments of the de novo peptides of the invention are preferred. in the K<sup>+</sup>(G/Sar)D sequence, G/Sar is preferably G. AA<sub>1</sub> and AA<sub>4</sub> are preferably Gly, Ala or Ser; n<sub>1</sub> is preferably 0-2, n<sub>4</sub> is preferably 1-2.

- 5 Preferred for AA<sub>2</sub> are neutral/nonpolar/aromatic amino acids, especially tryptophan and phenylalanine, particularly tryptophan, n<sub>2</sub> is preferably 1. X<sub>1</sub> and X<sub>2</sub> are preferably Cys, Mpr, or Pen (penicillamine) residues. Y<sub>1</sub> is preferably H, acetyl, or Gly; Y<sub>2</sub> is preferably -NH<sub>2</sub> or -A-NH<sub>2</sub>. Also preferred generally are C-terminal  
10 amidated forms of Y<sub>2</sub>.

- Thus, preferred embodiments of the PAI analogs include peptides of the following formulas. Although all of these are capable of provision in cyclic form through  
15 formation of disulfide linkages, these linkages are not specifically shown; other cyclic forms are noted by "cyclo."

#### Preferred Peptides

20

PAI 1: E-C-A-D-G-L-C-C-D-Q-C-R-F-L-K-K-G-T-V-  
C-R-V-A-K-G-D-W-N-D-D-T-C-T-G-Q-S-C-D-C-P-R-N-G-L-Y-G

PAI 2: E-E-P-C-A-T-G-P-C-C-R-R-C-K-F-K-R-A-G-  
K-V-C-R-V-A-K-G-D-W-N-N-D-Y-C-T-G-K-S-C-D-C-P-R-N-P-W-N-G

25

PAI 3: G-C-G-K-G-D-W-P-C-A-NH<sub>2</sub>;

PAI 4: G-C-K-G-D-W-P-C-A-NH<sub>2</sub>

PAI 5: C-G-K-G-D-W-P-C-NH<sub>2</sub>

PAI 7: C-K-G-D-W-C-A-NH<sub>2</sub>;

30

PAI 9: Mpr-K-G-D-Pen-NH<sub>2</sub>

35



	PAI 10:	C-K-G-D-W-P-C-NH <sub>2</sub>
	PAI 12:	C-K-G-D-Y-P-C-NH <sub>2</sub>
5	PAI 13:	C-K-G-D-F-P-C-NH <sub>2</sub>
	PAI 14:	C-K-G-D-L-P-C-NH <sub>2</sub>
	PAI 15:	C-K-G-D-V-P-C-NH <sub>2</sub>
	PAI 16:	C-K-G-D-Y(OMe)-P-C-NH <sub>2</sub>
10	PAI 17:	C-K-G-D-(2-Nal)-P-C-NH <sub>2</sub>
	PAI 18:	C-K-G-D-(Cha)-P-C-NH <sub>2</sub>
	PAI 19:	Mpr-K-G-D-W-P-C-NH <sub>2</sub>
	PAI 20:	Mpr-K-G-D-Y-P-C-NH <sub>2</sub>
15	PAI 21:	Mpr-K-G-D-F-P-C-NH <sub>2</sub>
	PAI 22:	Mpr-K-G-D-L-P-C-NH <sub>2</sub>
	PAI 23:	Mpr-K-G-D-V-P-C-NH <sub>2</sub>
	PAI 24:	Mpr-K-G-D-Y(OMe)-P-C-NH <sub>2</sub>
20	PAI 25:	Mpr-K-G-D-(2-Nal)-P-C-NH <sub>2</sub>
	PAI 26:	Mpr-K-G-D-(Cha)-P-C-NH <sub>2</sub>
	PAI 27:	cyclo(G-K-G-D-W-P)
	PAI 28:	cyclo(A-K-G-D-W-P)
	PAI 29:	cyclo(D-Ala-K-G-D-W-P)
25	PAI 30:	cyclo(F-K-G-D-W-P)
	PAI 31:	cyclo(beta-Ala-K-G-D-W-P)
	PAI 32:	cyclo(gamma-Abu-K-G-D-W-P)
	PAI 33:	cyclo(R-K-G-D-W-P)
30	PAI 34:	C-K-G-D-W-G-C-NH <sub>2</sub>
	PAI 37:	C-K-A-D-W-P-C-NH <sub>2</sub>
	PAI 39:	C-K-G-D-W-(Sar)-C-NH <sub>2</sub>
35	PAI 41:	C-K-G-D-I-P-C-NH <sub>2</sub>

	PAI 42:	C-K-G-D-(4-Cl-Phe)-P-NH <sub>2</sub>
	PAI 43:	C-K-(Sar)-D-W-P-C-NH <sub>2</sub>
5	PAI 44:	C-K-G-D-(4-NO <sub>2</sub> -Phe)-P-C-NH <sub>2</sub>
	PAI 47:	Acetyl-C-K-G-D-W-P-C-NH <sub>2</sub>
	PAI 48:	Mpr-K-G-D-W(Formyl)-P-C-NH <sub>2</sub>
	PAI 49:	Mvl-K-G-D-W-P-C-NH <sub>2</sub>
10	PAI 51:	Mpr-K-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 52:	Mpr-K-G-D-W-P-Pen <sup>†</sup> -NH <sub>2</sub>
	PAI 54:	Mpr-K-G-D <sup>†</sup> -W-P-Pen-NH <sub>2</sub>
	PAI 55:	Mpr-K-G-D-W-(Thz)-C-NH <sub>2</sub>
15	PAI 56:	Mpr-K-G-D-H(2,4-DNP)-P-C-NH <sub>2</sub>
	PAI 57:	Mpr-K-G-D-(2-Nal)-P-Pen-NH <sub>2</sub>
	PAI 58:	Mvl-K-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 59:	Mpr-K-G-D-W-(Pip)-Pen-NH <sub>2</sub>
20	PAI 60:	Mpr-(Har)-G-D-W-P-C-NH <sub>2</sub>
	PAI 61:	Mpr-K-G-D-W-P-C <sup>†</sup> -NH <sub>2</sub>
	PAI 62:	Mpr-K <sup>†</sup> -G-D-W-P-Pen-NH <sub>2</sub>
	PAI 63:	Mpr-(Har)-G-D-W-P-Pen-NH <sub>2</sub>
25	PAI 64:	Mpr-(Acetimidyl-Lys)-G-D-W-P-C-NH <sub>2</sub>
	PAI 65:	Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 66:	Mpr-(N <sup>G</sup> ,N <sup>G'</sup> -ethylene-Har)-G-D-W-P-C-NH <sub>2</sub>
	PAI 67:	Mpr-(N <sup>G</sup> ,N <sup>G'</sup> -ethylene-Har)-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 68:	Mpr-Har-Sar-D-W-P-C-NH <sub>2</sub>
30	PAI 69:	Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 70:	Mpr-(Phenylimidyl-Lys)-G-D-W-P-C-NH <sub>2</sub>
	PAI 71:	Mpr-Har-Sar-D-W-P-PenNH <sub>2</sub>
35	PAI 72:	Mpr-(Phenylimidyl-Lys)-G-D-W-P-PenNH <sub>2</sub>

SUBSTITUTE SHEET

-25-

	PAI 73	Mpr-Har-G-D-W- (3,4-dehydro-Pro) -C-NH <sub>2</sub>
	PAI 74	Mpr-Har-G-D-Pen-NH <sub>2</sub>
	PAI 75	Mpr- (Phenylimidyl-Lys) -G-D-Pen-NH <sub>2</sub>
5	PAI 80	Mpr-P-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 81	Mpr-G-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 82	Mpr-A-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 83	Mpr-Aib-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 84	Mpr- (N-Me-Arg) -Har-G-D-W-P-C-NH <sub>2</sub>
10	PAI 85	Mpr- (N-Me-Ser) -Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 86	Mpr-A <sup>†</sup> -Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 87	Mpr- (β-Ala) -Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 88	Mpr- (N-Me-Leu) -Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 89	Mpr- (N-Me-Ala) -Har-G-D-W-P-C-NH <sub>2</sub>
15	PAI 90	Mpr-Sar-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 91	Mpr-V-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 92	Mpr-S-Har-G-D-W-P-C-NH <sub>2</sub>
	PAI 93	Mpr-Har-G-D-W-P-A-C-NH <sub>2</sub>
	PAI 94	Mpr-Har-G-D-W-P- (N-Me-Ala) -C-NH <sub>2</sub>
20	PAI 95	Mpr-Har-G-D-W-P-G-C-NH <sub>2</sub>
	PAI 96	Mpr-Har-G-D-W-P-A <sup>†</sup> -C-NH <sub>2</sub>
	PAI 97	Mpr-Har-G-D-W-P-P-C-NH <sub>2</sub>
	PAI 98	Mpr-Har-G-D-W-P- (Sar) -C-NH <sub>2</sub>
	PAI 99	Mpr-Har-G-D-W-P- (Aib) -NH <sub>2</sub>
25	PAI 100	Mpr-A- (Har) -G-D-W-P-Pen-NH <sub>2</sub>
	PAI 101	Mpr-A-K-G-D-W-P-Pen-NH <sub>2</sub>
	PAI 102	Mpr-D- (Har) -G-D-W-P-Pen-NH <sub>2</sub>

30

35

SUBSTITUTE SHEET

Particularly preferred are peptides of the formulas

- 5 PAI 3: G-C-G-K-G-D-W-P-C-A-NH<sub>2</sub>;  
 PAI 4: G-C-K-G-D-W-P-C-A-NH<sub>2</sub>;  
 PAI 5: C-G-K-G-D-W-P-C-NH<sub>2</sub>;  
 PAI 9: Mpr-K-G-D-Pen-NH<sub>2</sub>; and  
 PAI 10: C-K-G-D-W-P-C-NH<sub>2</sub>  
 10 PAI 12: C-K-G-D-Y-P-C-NH<sub>2</sub>  
 PAI 13: C-K-G-D-F-P-C-NH<sub>2</sub>  
 PAI 19: Mpr-K-G-D-W-P-C-NH<sub>2</sub>  
 PAI 25: Mpr-K-G-D-(2-Nal)-P-C-NH<sub>2</sub>  
 PAI 34: C-K-G-D-W-G-C-NH<sub>2</sub>  
 15 PAI 39: C-K-G-D-W-(Sar)-C-NH<sub>2</sub>  
 PAI 42: C-K-G-D-(4-Cl-Phe)-P-NH<sub>2</sub>  
 PAI 43: C-K-(Sar)-D-W-P-C-NH<sub>2</sub>  
 PAI 44: C-K-G-D-(4-NO<sub>2</sub>-Phe)-P-C-NH<sub>2</sub>  
 20 PAI 47: Acetyl-C-K-G-D-W-P-C-NH<sub>2</sub>  
 PAI 48: Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>  
 PAI 49: Mvl-K-G-D-W-P-C-NH<sub>2</sub>  
 PAI 51: Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>  
 25 PAI 52: Mpr-K-G-D-W-P-(D-Pen)-NH<sub>2</sub>  
 PAI 55: Mpr-K-G-D-W-(Thz)-C-NH<sub>2</sub>  
 PAI 56: Mpr-K-G-D-H(2,4-DNP)-P-C-NH<sub>2</sub>  
 PAI 57: Mpr-K-G-D-(2-Nal)-P-Pen-NH<sub>2</sub>  
 30 PAI 58: Mvl-K-G-D-W-P-Pen-NH<sub>2</sub>  
 PAI 59: Mpr-K-G-D-W-(Pip)-Pen-NH<sub>2</sub>  
 PAI 60: Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 PAI 61: Mpr-K-G-D-W-P-C<sup>†</sup>-NH<sub>2</sub>  
 35

- PAI 62: Mpr-K<sup>†</sup>-G-D-W-P-Pen-NH<sub>2</sub>  
 PAI 63: Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>  
 5 PAI 64: Mpr-(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 PAI 65: Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 PAI 66: Mpr (N<sup>G</sup>,N<sup>G'</sup>-ethylene-Har)-G-D-W-P-C-NH<sub>2</sub>  
 PAI 67: Mpr (N<sup>G</sup>,N<sup>G'</sup>-ethylene-Har)-G-D-W-P-Pen-NH<sub>2</sub>  
 10 PAI 68: Mpr-Har-Sar-D-W-P-C-NH<sub>2</sub>  
 PAI 69: Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 PAI 70: Mpr-(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 PAI 71: Mpr-Har-Sar-D-W-P-PenNH<sub>2</sub>  
 PAI 72: Mpr-(Phenylimidyl-Lys)-G-D-W-P-PenNH<sub>2</sub>  
 15 PAI 73: Mpr-Har-G-D-W-(3,4-dehydro-Pro)-C-NH<sub>2</sub>

#### Chemical Synthesis of the Invention Peptides

- 20 Polypeptides within the scope of the present invention can be synthesized chemically by means well known in the art such as, e.g., solid-phase peptide synthesis. The synthesis is commenced from the carboxy-terminal end of the peptide using an alpha-amino  
 25 protected amino acid. t-Butyloxycarbonyl (Boc) protective groups can be used for all amino groups even though other protective groups such as fluorenylmethyloxycarbonyl (Fmoc), are suitable. For example, Boc-Gly-OH, Boc-Ala,OH, Boc-His (Tos)-OH, (i.e.,  
 30 selected carboxy-terminal amino acids) can be esterified to chloromethylated polystyrene resin supports, p-methyl benzhydrylamine (pMBHA) or PAM resins. The polystyrene resin support is preferably a copolymer of styrene with about 0.5 to 2% divinyl benzene as a cross-linking agent  
 35 which causes the polystyrene polymer to be completely

**SUBSTITUTE SHEET**

-28-

insoluble in certain organic solvents. See Stewart, et al., Solid-Phase Peptide Synthesis (1969) W.H. Freeman Co., San Francisco and Merrifield J Am Chem Soc (1963) 85:2149-2154. These and other methods of peptide synthesis are also exemplified by U.S. Patent Nos. 3,862,925, 3,842,067, 3,972,859, and 4,105,602.

The synthesis can use manual synthesis techniques or automatically employ, for example, an Applied BioSystems 430A or 431A Peptide Synthesizer (Foster City, California) following the instructions provided in the instruction manual supplied by the manufacturer. Cleavage of the peptides from the resin can be performed using the "low-high" HF deprotection protocols as described in Lu, G.-S., et al., Int J Peptide & Protein Res (1987) 29:545-557. Refolding of analogs of the snake venom PAIs can be performed using the procedure outlined in Garsky, V., et al., Proc natl Acad Sci USA (1989) 86:4022-4026 which describes the solid-phase synthesis of echistatin.

The cyclic peptides of this invention which do not have disulfide bonds can be conveniently prepared by a combination of solid phase synthesis and formation of the cyclic ring structure in solution using the general methods as outlined in U.S. Patent 4,612,366 to Nutt. Thus, linear peptides prepared on standard Merrifield resin can be cleaved from the resin with hydrazine, followed by cyclization of the corresponding azide to form the cyclic peptides.

It will be readily appreciated by those having ordinary skill in the art of peptide synthesis that the intermediates which are constructed in accordance with the present disclosure during the course of synthesizing the present analog compounds are themselves novel and useful compounds and are thus within the scope of the invention.

SUBSTITUTE SHEET

Recombinant Production

Alternatively, selected polypeptides of the present invention can be produced by expression of recombinant DNA constructs prepared in accordance with well-known methods. Such production can be desirable to provide large quantities or alternative embodiments of such compounds. Since the peptide sequences are relatively short, recombinant production is facilitated; however, production by recombinant means is particularly preferred over standard solid phase peptide synthesis for peptides of at least 8 amino acid residues.

The DNA encoding the sequenced polypeptide, such as a PAI, is preferably prepared using commercially available nucleic acid synthesis methods. Methods to construct expression systems for production of PAI in recombinant hosts are also generally known in the art.

Expression can be effected in either procaryotic or eucaryotic hosts. Procaryotes most frequently are represented by various strains of E. coli. However, other microbial strains may also be used, such as bacilli, for example Bacillus subtilis, various species of Pseudomonas, or other bacterial strains. In such procaryotic systems, plasmid vectors which contain replication sites and control sequences derived from a species compatible with the host are used. For example, a workhorse vector for E. coli is pBR322 and its derivatives. Commonly used procaryotic control sequences, which contain promoters for transcription initiation, optionally with an operator, along with ribosome binding-site sequences, include such commonly used promoters as the beta-lactamase (penicillinase) and lactose (lac) promoter systems, the tryptophan (trp) promoter system, and the lambda-derived  $P_L$  promoter and N-gene ribosome binding site. However, any available promoter system compatible with procaryotes can be used.

-30-

Expression systems useful in eucaryotic hosts comprise promoters derived from appropriate eucaryotic genes. A class of promoters useful in yeast, for example, includes promoters for synthesis of glycolytic enzymes, e.g., those for 3-phosphoglycerate kinase. Other yeast promoters include those from the enolase gene or the Leu2 gene obtained from YEpl3.

Suitable mammalian promoters include the early and late promoters from SV40 or other viral promoters such as those derived from polyoma, adenovirus II, bovine papilloma virus or avian sarcoma viruses. Suitable viral and mammalian enhancers are cited above. In the event plant cells are used as an expression system, the nopaline synthesis promoter, for example, is appropriate.

The expression systems are constructed using well-known restriction and ligation techniques and transformed into appropriate hosts.

Transformation is done using standard techniques appropriate to such cells. The cells containing the expression systems are cultured under conditions appropriate for production of the polypeptide, e.g., PAI and the polypeptide, e.g., PAI, is then recovered and purified.

## 25 Antibodies

The availability of the purified polypeptide, e.g., PAI, of the invention also permits the production of antibodies specifically immunoreactive with these forms of the active peptide.

For example, the compositions containing purified PAI isolated from snake venom or otherwise synthesized can be used to stimulate the production of antibodies which immunoreact with the PAI peptide. Standard immunization protocols involving administering PAI to various vertebrates, such as rabbits, rats, mice, sheep,

SUBSTITUTE SHEET



-31-

and chickens result in antisera which are immunoreactive with the purified peptide. Polypeptide, e.g., PAI, may be advantageously conjugated to a suitable antigenically neutral carrier, such as an appropriate serum albumin or keyhole limpet hemocyanin, in order to enhance immunogenicity. In addition, the free peptide can be injected with methylated BSA as an alternative to conjugation. Furthermore, the antibody-secreting cells of the immunized mammal can be immortalized to generate monoclonal antibody panels which can then be screened for reactivity with the polypeptide, such as PAI.

The resulting polyclonal or monoclonal antibody preparations are useful in assays for levels of the corresponding polypeptide, e.g., PAI, in biological samples using standard immunoassay procedures.

#### Active PAI Assay

The identification of snake venom starting material which contains active PAI, and which PAI has known specificity, is made possible by the PAI assay. The PAI assay rests on the observation that compounds which block the binding of fibrinogen to the GP IIb-IIIa complex in vitro also are capable of inhibiting thrombin or ADP-induced aggregation of human platelets and the formation of platelet-thrombi in vivo. This observation provides the basis for obtaining potent PAIs by evaluating the ability of test materials to disrupt fibrinogen-GP IIb-IIIa interactions.

In the assay, GP IIb-IIIa, prepared in purified form, for example as described by Fitzgerald, L.A., et al., Anal Biochem (1985) 151:169-177, incorporated herein by reference, is coated onto a solid support such as beads, test tubes, or microtiter plates. The coated support is then contacted with fibrinogen and with the test material and incubated for a sufficient time to

SUBSTITUTE SHEET

-32-

5 permit maximal binding of fibrinogen to the immobilized GP IIb-IIIa. Fibrinogen is typically provided at a concentration of about 5-50 nM and the test material can, if desired, be added at a series of dilutions. Typical incubations are 2-4 hr at 35°C, the time and temperature being interdependent.

10 After incubation, the solution containing the fibrinogen and test material is removed and the level of binding of fibrinogen measured by quantitating bound fibrinogen to GP IIb-IIIa. Any suitable means of detection can be used, but it is convenient to employ labeled fibrinogen, for example using radioactive, fluorescent or biotinylated labels. Such methods are well known and need not be elaborated here.

15 Assessment of the results is aided by employing a control sample, usually identical to the test sample except that the test substance is absent. In this case, percent inhibition may be calculated using the basis, so that

20 
$$\% \text{ inhibition} = \frac{\text{control} - \text{test}}{\text{control}} \times 100.$$

Other measures of inhibition effectiveness, such as  $IC_{50}$ , can also be used.

25 The PAI assay systems further include characterization of the PAI specificity by binding inhibition assays identical to that above but substituting other adhesive proteins for Fg and other receptors for GP IIb-IIIa. In particular, inhibition of the binding of vitronectin to the vitronectin receptor; fibronectin to the fibronectin receptor; fibronectin to GP IIb-IIIa and fibrinogen and/or vWF to GP IIb-IIIa can be assessed. The adhesive protein and receptors for these assays are available in the art.

35

SUBSTITUTE SHEET

-33-

Other Assays

In addition to the above plate assays, other assays for platelet aggregation inhibition activity and related activities are also available, as set forth above. In summary, a list of commonly employed assays is as follows:

1. The plate assays utilizing specific receptors described in the previous paragraphs;
2. Standard assays directly applied to platelet aggregation, such as those described by Gann, Z.-R., et al., J Biol Chem (1988) 263:19827-19832; Huang, T.F., et al., J Biol Chem (1987) 262:16157-16163; Biochemistry (1989) 28:661-666, cited above and incorporated herein by reference;
3. An in vivo thrombosis model in dogs as described by Folts, J.D., et al., Circulation (1976) 54:365; and
4. Effect on cell adhesion using S35 methionine-labeled cells as described hereinbelow in Example 10.

Administration and Utility of PAIs

The PAIs of the invention are useful therapeutically to prevent thrombus formation. Indications appropriate to such treatment include, without limitation, atherosclerosis and arteriosclerosis, acute myocardial infarction, chronic unstable angina, transient ischemic attacks and strokes, peripheral vascular disease, arterial thrombosis, preeclampsia, embolism, restenosis and/or thrombosis following angioplasty, carotid endarterectomy, anastomosis of vascular grafts, and chronic cardiovascular devices (e.g., in-dwelling catheters or shunts "extracorporeal circulating devices"). These syndromes represent a variety of stenotic and occlusive vascular disorders thought to be initiated by platelet activation on vessel walls.

SUBSTITUTE SHEET

-34-

The PAIs can be used for prevention or abortion of arterial thrombus formation, in unstable angina and arterial emboli or thrombosis, as well as treatment or preventions of myocardial infarction (MI) and mural thrombus formation post MI. For brain-related disorders, treatment or prevention of transient ischemic attack and treatment of thrombotic stroke or stroke-in-evolution are included.

The PAIs can also be used for prevention of platelet aggregation, embolization, or consumption in extracorporeal circulations, including improving renal dialysis, cardiopulmonary bypasses, hemoperfusions, and plasmapheresis.

PAIs prevent platelet aggregation, embolization, or consumption associated with intravascular devices, and administration results in improved utility of intraaortic balloon pumps, ventricular assist devices, and arterial catheters.

The PAIs will also be useful in treatment or prevention of venous thrombosis as in deep venous thrombosis, IVC, renal vein or portal vein thrombosis, and pulmonary venous thrombosis.

Various disorders involving platelet consumption, such as thrombotic thrombocytopenic purpura are also treatable.

In addition, the PAIs of the present invention can be used in numerous nontherapeutic applications where inhibiting platelet aggregation is desired. For example, improved platelet and whole blood storage can be obtained by adding sufficient quantities of the peptides, the amount of which will vary depending upon the length of proposed storage time, the conditions of storage, the ultimate use of the stored material, etc.

The PAI dosage can range broadly depending upon the desired affects and the therapeutic setting. Typically,

SUBSTITUTE SHEET

-35-

dosages can be a 0.5 mg/ml dosage form, which is a direct infusion pre-mix is a bolus followed by an infusion. The bolus is about 10 to about 500, preferably about 30-300  $\mu\text{g/Kg}$  (about 1-40 mg, preferably from about 2-20 mg for a 70Kg patient) and the infusion is 0.02-2  $\mu\text{g/Kg/min}$ , but probably 0.5-1.0  $\mu\text{g/Kg/min}$  (50-100 mg/day for a 70Kg patient). Administration is preferably parenteral, such as intravenous on a daily basis for up to a week or as much as one or two months or more, all of which will vary with the peptide's size. If the peptides are sufficiently small (e.g., less than about 8-10 amino acid residues) other routes of administration can be utilized, such as intranasally, sublingually, or the like.

Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions.

The invention also includes novel therapeutic compositions consisting essentially of an injectable biologically active substantially pure polypeptide as a liquid solution in a citrate buffer, said solution having a pH of from about 5.0 to about 5.5. The compositions are capable of remaining very stable at about 4°C, are still stable even at about 50°C and have improved stability at about 70°C for 7, 34 or 49 days prior to injection as compared to compositions having a pH greater than about 5.5. The compositions of the invention also remain stable and injectable at about -15° C to about 30° C for at least 18 months.

The therapeutic compositions contain any injectable biologically active substantially pure polypeptide as discussed above. Preferably the substantially pure polypeptide is biologically active for inhibiting thrombus formation, preventing platelet loss during extracorporeal circulation of blood or for treating a

SUBSTITUTE SHEET

-36-

patient suspected of having a platelet-associated ischemic syndrome. In one preferred embodiment, the polypeptide is a cyclic polypeptide containing up to 10 amino acid residues, and preferably at least one

5 disulfide bond, such as Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>, Mvl-K-G-D-W-P-C-NH<sub>2</sub>, Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>, Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>, Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>, Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>, Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>, Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>,

10 Mpr(Phenylimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>, Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub>, Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub>, Mpr-K-G-D-W-P-C-NH<sub>2</sub>, or cyclic forms thereof. Preferably, the polypeptide is Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub>

• acetate or a cyclic form thereof.

15 In another aspect, the invention includes novel PAI in isolated form which is identified in, and can be isolated form, active snake venom according to the methods of the invention. In particular, the invention relates to PAI, in isolated form, which can be isolated

20 from Echis colorata, Eristicophis macmahonii; A. hypnale, A. acutus, A. piscivorous leucostoma, A. piscivorous conanti; Bothrops asper; Bothrops cotiara, B. jararaca, B. jararacussu, B. lansbergi, B. medusa, B. nasuta, B. neuwiedi, B. pradoi, B. schlegli; Crotalus atrox, C. basilicus, C. cerastes cerastes, C. durissus durissus, C. durissus totonatacus, C. horridus horridus, C. molossus mollossus, C. ruber ruber, C. viridis cereberus, Crotalus v. helleri, Crotalus v. lutosus, Crotalus v. oreganus, Crotalus v. viridis; Lachesis mutas; Sistrurus catenatus

25 tergeminus, and Sistrurus milarus barbouri.

Preferred are PAIs in isolated form prepared from, or having the amino acid sequences of, those obtained from Eristicophis macmahonii (eristicophin); Bothrops cotiara (cotiarin); B. jararacussu; Crotalus atrox

35 (crotatoxin); C. basilicus (basilicin); C. cerastes

-37-

cerastes (cerastin); C. durissus totonatacus (durissin);  
C. durissus durissus (durissin); C. h. horridus  
(horridin); Crotalus m. mollossus (molossin); C. ruber  
ruber (ruberin); Crotalus viridis lutosus (lutosin); C.  
5 v. viridis (viridin); Crotalus v. oreganus (oreganin);  
Crotalus v. helleri; Lachesis mutas (lachesin); Sistrurus  
catenatus tergeminus (tergeminin); and S. milarus  
barbouri (barbourin).

Especially preferred are eristocophin, cotiarin,  
10 crotatroxin, cerastin, durissin, horridin, ruberin,  
lachesin, basilicin, lutosin, molossin, oreganin,  
viridin, tergeminin and barbourin.

The invention also includes peptides of the amino  
acid sequences as described above which are truncated  
15 and/or modified forms of the naturally occurring peptides  
and/or have one or more peptide linkages replaced by  
alternate linkages such as -CH<sub>2</sub>NH- or -CH<sub>2</sub>CH<sub>2</sub>-.

In another aspect, the invention relates to PAI in  
isolated form which can be prepared from active snake  
20 venom identified by the method of the invention, and  
shown to specifically inhibit the binding fibrinogen (Fg)  
and/or von Willebrand Factor (vWF) to GP IIb-IIIa, and  
their truncated and/or modified forms.

In still another aspect, the invention relates to  
25 PAI of snake venom in isolated form wherein the sequence  
responsible for binding to the adhesive protein receptor  
includes the sequence KGD.

In another aspect, the invention is directed to a  
group of peptides or peptide-related compounds in general  
30 which are platelet aggregation inhibitors that are  
capable of inhibiting binding of Fg or vWF to GP IIb-IIIa  
at a substantially higher potency than that at which they  
inhibit binding of vitronectin to vitronectin receptor or  
fibronectin to fibronectin receptor. These peptides are  
35 characterized by having the binding sequence K<sup>\*</sup>GDX in

-38-

place of the RGD<sub>X</sub> binding sequence which is found in other PAI proteins. K<sup>\*</sup> is a substituted or unsubstituted lysyl-derived residue of the formula R<sup>1</sup><sub>2</sub>N(CH<sub>2</sub>)<sub>4</sub>CH(NH)CO- wherein each R<sup>1</sup> is independently H or a substituent which is sufficiently electron donating so as to not destroy the basicity of the adjacent nitrogen, and wherein one or two of the methylene residues may optionally be substituted by O or S, as described below. The barbourin PAI isolated from S. milarus barbouri is one illustration of this series of peptides. However, shorter forms of this peptide can also be used, as well as analogous sequences which also contain 1-10 amino acid residue modifications elsewhere in the peptide chain, and/or replacement of peptide linkages with alternate linkages. Other illustrative embodiments include isolated PAI peptides having a negative RGD<sub>X</sub> sequence wherein this is replaced by K<sup>\*</sup>GD<sub>X</sub>. As in the case of barbourin, these isolated PAI may be otherwise in native form, or may be truncated and/or may contain 1-10 amino acid residue substitutions or deletions, and/or may have non-peptide linkages substituted for peptide linkages.

Another group of compounds which falls within the scope of the invention is that wherein the foregoing compounds are as described, except that the glycyl residue in the RGD or K<sup>\*</sup>GD sequence is replaced by a sarcosyl residue. This class of compounds retains the potency and specificity of the related RGD or K<sup>\*</sup>GD-containing peptides.

The platelet aggregation inhibitors (PAI) of the invention include low molecular weight peptides which can be prepared in isolated form, as described below, from snake venom which has been identified as "active," i.e., has been found to contain PAI using the method of the invention, which is described hereinbelow.

SUBSTITUTE SHEET



-39-

The invention method permits ready identification and characterization of the presence of an effective PAI in snake venom which selectively inhibits binding to GP IIb-IIIa as opposed to other integrins as, for example, the vitronectin receptor and the fibronectin receptor. Upon such identification, and, optionally and optimally, characterization, the PAI can be isolated and purified using a variety of standard techniques illustrated herein and disclosed in the art. For example, a combination of separation based on molecular weight (typically recovery of substances of <10kd), ion exchange chromatography, and reverse phase HPLC can be used. Other techniques can also be employed, but a workable procedure applicable to PAI from any active snake venom is as follows:

About 10-1000 mg venom is dissolved in dilute acetic acid and applied to a sizing column, such as Sephadex G-50, and eluted in the same solvent. Fractions are assayed for activity using the Fg/GP IIb-IIIa binding assay of the invention, a standard platelet aggregation assay (PAA) or any similar assay relying on the adhesive protein binding activity of GP IIb-IIIa. Alternatively, the <10kd fraction of the fraction of the venom can be recovered using ultrafiltration and similarly assayed.

The low MW fraction isolated by either procedure is then loaded onto a preparative C-18 HPLC column, such as a C-18 Delta Pak reverse phase HPLC column, available from Waters, preequilibrated in 0.1% trifluoroacetic acid (TFA)/8% acetonitrile. The adsorbed PAI is then eluted using a gradient of 8%-60% acetonitrile in 0.1% TFA. The slope of the gradient and flow rate are optimized using routine procedures. Active fractions are determined by PAA or by the disclosed receptor binding method. The active fractions are then pooled, concentrated, and tested for homogeneity using analytical HPLC or SDS-PAGE.

SUBSTITUTE SHEET

-40-

Further reverse-phase HPLC gradient purification is applied until the recovered PAI is homogenous.

5 It is understood that the isolated PAI of determined sequence can, when synthesized in vitro, be modified by sequence alterations which do not destroy activity. In general, these modified forms will differ from the native forms by 1-10, preferably 1-4, amino acid substitutions or will be truncated forms. In addition, one or more peptide linkages may be replaced by alternate linkages as described herein. A particularly preferred substitution is replacement of RGD by K\*GD to confer GP IIb-IIIa specificity as described herein.

The PAI of Sistrurus m. barbouri has been purified to homogeneity and sequenced, and termed "barbourin".  
15 Unlike the adhesive proteins for GP IIb-IIIa so far identified and the peptides from snake venoms that block GP IIb-IIIa function, barbourin does not contain the standard Arg-Gly-Asp sequence of the adhesive proteins known in the art. The apparent binding sequence in  
20 barbourin is Lys-Gly-Asp-(Trp). The presence of the KGD sequence in the apparent binding region of this peptide is especially surprising in view of the observation that replacement of Lys for Arg in small synthetic peptides based on the RDGX sequence greatly decreases the ability  
25 of these peptides to bind to integrin receptors (Pierschbacher et al., Proc Natl Acad Sci (USA) (1984) 81:5985-5988; Williams et al., Thromb Res (1987) 46:457-471); Huang et al., J. Biol Chem (1987) 262:16157-16163. It is thought that this substitution may in part be  
30 responsible for the specificity of the barbourin peptide to inhibit Fg and vWF binding to GP IIb-IIIa, versus, for example, inhibition of vitronectin binding to the vitronectin receptor.

35

SUBSTITUTE SHEET

-41-

Examples

The invention is illustrated by the following examples which should not be regarded as limiting the invention in any way.

5

Example 1 Preparation of Stabilized Composition

Citrate buffer solution is prepared by mixing citric acid in a volumetric flask to give a final citrate concentration of 25 mM with Nanopure water and adding  
10 NaOH to adjust the pH.

To the above citrate buffer solution was added 200 ng of greater than 95% pure cyclic form of Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate polypeptide (I) to give a solution concentration of 2  
15 mg/ml when adjusted to the desired pH.

Stability Determination

One ml aliquots of the above solution of polypeptide in citrate buffer were sealed in borosilicate glass ampules under sterile conditions and stored at a constant  
20 temperature. Test sample ampules from each pH and temperature condition 4°C, 50°C and 70°C were removed periodically, visually assessed and then analyzed separately by HPLC using a mobile phase comprising acetonitrile (MeCN) and 0.1% trifluoroacetic (TFA) in  
25 water. This stability was evaluated by conventional assay methods applicable to purity, weight or size of polypeptides. These included not only visual evaluations, such as discoloration, transparency and precipitation, and included assays normally applied to  
30 separate polypeptides from each other and from other materials by reversed phase high performance liquid chromatography (HPLC).

Results of these experiments are set forth in Table 1 in which the stability is expressed in terms of the  
35 percentage of main peak of original polypeptide found in

SUBSTITUTE SHEET

-42-

the sample analyzed by HPLC. The composition was also found to be normal on visual evaluation.

5

10

15

20

25

30

35

**SUBSTITUTE SHEET**

-43-

Table 1  
Peptide Stability in Citrate Buffer Composition

	4°C			50°C			70°C		
	pH=5.0	5.5	6.0	pH=5.0	5.5	6.0	pH=5.0	5.5	6.0
Time (min)									
21.0	100	100	100	99	100	98	99	96	94
45.5	100	99	98	97	95	94	91	88	83
167.5	100	99	98	98	99	97	80	80	68
503.5	98	99	96	94	92	89	47	51	33
837.5	100	101	100	95	94	88	29	36	20
1173.5	100	101	100	92	92	83	17	22	13

SUBSTITUTE SHEET

-44-

Results of these experiments demonstrate that liquid solutions of I in citrate buffer having a pH of 5.00 and 5.50 were unexpectedly very stable at about 4°C, still stable at up to about 50°C and much more stable at 70°C in comparison to citrate buffer solutions at pH 6.0.

Example 2 Preparation of Stabilized Compositions

A solution was formed in water of a peptide of the invention (greater than 95% pure cyclic form of Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate polypeptide (I)). To this solution was added about 1.6 mg/ml of citric acid monohydrate followed by about 5.1 mg/ml of sodium citrate dihydrate to give a final citrate solution concentration of 25 nM in water for injection and a final polypeptide concentration of about 2 mg/ml.

Stability Determination

20 ml aliquots of the above solution were placed into Type I glass vials and sealed with rubber stoppers and aluminum closures under sterile conditions and stored at a constant temperature. Test sample vials from each temperature condition, -15°C, 5°C, 30°C and 45°C were removed periodically and individually assessed and analyzed for stability by HPLC and visual evaluation as described in Example 1. Results of these experiments are set forth in Table 2 below.

-45-

Table 2  
Peptide Stability in 5.5 pH Citrate Buffer Composition

Time (days)	5°C	-15°C	30°C	45°C
0	98.8	-	-	-
45	-	98.9	97.4	96.1
76	98.6	-	-	-
95	-	99.0	96.4	94.1
126	98.5	-	-	-
191	-	99.1	95.8	92.4
222	98.5	-	-	-
270	-	98.7	95.1	89.8
301	98.2	-	-	-
368	-	98.6	94.5	86.5
399	98.2	-	-	-
541	-	98.9	93.6	76.1
572	98.2	-	-	-

\* After one month at 5°C, these samples were placed at the assigned temperature.

SUBSTITUTE SHEET

-46-

Results of these experiments demonstrated that liquid solutions of the polypeptide of the invention in citrate buffer having a pH of 5.25 were very stable from about -15°C to about 30°C. The composition was also found to be normal by visual evaluation.

### Example 3 Stability Determination

Following procedures similar to those described in Examples 1 and 2 above, Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub> was dissolved in 25 mM citrate buffer at pH 5.25 at a concentration of about 1 mg/mL and stored in test sample vials at 5°C, 30°C and 45°C for up to 90 days. Test sample vials from each temperature condition were removed periodically and analyzed for stability by HPLC and visual inspection as described in Example 1. Results of these experiments are set forth in Table 3 below.

Table 3

	<u>Time (Days)</u>	<u>5°C</u>	<u>30°C</u>	<u>45°C</u>
20	0	99.6	-	-
	29	98.1	97.3	95.4
	60	98.3	97.3	92.6
	90	97.3	96.5	90.6

Results of these experiments demonstrated that liquid solutions of the polypeptide of the invention having a pH of 5.25 were very stable from about 5°C to about 45°C. The composition was also found to be normal on visual evaluation.

### Example 4

#### Preparation of Analog #1

[E<sup>28</sup>L<sup>41</sup>C<sup>64</sup>]barbourin (28-73):

E-C-A-D-G-L-C-C-D-O-C-R-F-L-K-K-G-T-V-C-R-V-

A-K-G-D-W-N-D-D-T-C-T-G-O-S-C-D-C-P-R-N-G-L-Y-G



-47-

One-half mmol of PAM-Gly resin (0.6 meq/g, Applied Biosystems, Foster City, CA) was subjected to Procedure A with the required amino acids (introduced in order). The Boc-protected amino acids had the following side-chain  
5 protection: Arg(Tos), Asp(OcHex), Cys(4-MeBzl), Glu(OcHex), Lys(Cl-Z), Thr(OBzl), Trp(CHO), and Tyr(Br-Z). Following assembly of the completed protected peptide-resin chain, the amino terminal Boc- group was removed with TFA and the resin dried as its TFA-salt  
10 form. The resin (1.3 g) was subjected to "low-high" HF deprotection protocols followed by removal of HF "in vacuo." The dried peptide-resin mixture was transferred to a fretted funnel (coarse) with ethyl ether and was washed several times with alternate washes of ether and  
15 chloroform to remove most of the organic protecting groups and scavengers used in the deprotection.

The Peptide mixture was transferred to 2 L of 0.4% acetic acid and the pH adjusted to 7.99 with concentrated  $\text{NH}_4\text{OH}$ . The resin was filtered from this solution and the  
20 solution allowed to sit at 4°C without stirring for 20 hr. This was followed by warming the solution to room temperature and storing for 3 days again without stirring. Precipitated material was removed by filtration and the supernatant pH adjusted to 3.0 with  
25 acetic acid and lyophilized.

The crude material was dissolved in 8.0 ml of 0.5M acetic acid and loaded onto a Sephadex G-50 fine column (2.5 x 100 cm) equilibrated with 0.5M acetic acid. The  
30 column was run at 20 ml/hr and fractions (4 ml) were collected into polypropylene tubes. Aliquots of fractions were dried, resuspended in water and tested for platelet aggregation inhibitory activity as previously described. Active fractions (71-90) were pooled and  
35 lyophilized.

SUBSTITUTE SHEET

-48-

Dried material (66 mg) was redissolved in 2.0 ml of 0.1M acetic acid and loaded onto the Waters Preparative C-18 column equilibrated with 8% acetonitrile containing 0.1% TFA. A gradient running from 8% acetonitrile to 20% in 10 minutes followed by a slow gradient to 30% acetonitrile in 40 min was performed. The column was eluted at 18 ml/min and fractions (12 sec) were collected into polypropylene tubes. Fractions were concentrated on a Speed-Vac concentrator to 1.0 ml volume and 10 ul aliquots were tested in the platelet aggregation assay.

Active fractions (29-32) were individually lyophilized and analyzed on the analytical C-18 HPLC column with an 8-30% acetonitrile gradient. Fractions 29 and 30 were pooled and loaded onto the analytical column in 1.0 ml of 0.5% TFA. The major peak was collected manually and lyophilized to yield 1.6 mg of pure peptide.

Amino acid analysis of this material confirmed the identity of the peptide. Assay of this material for its ability to inhibit the binding of fibrinogen to GP IIb-IIIa and vitronectin demonstrated that the high affinity of this analog for GP IIb-IIIa and the relative lack of affinity for VnR at concentrations up to 1 uM.

#### Example 5

25 Preparation of Analog #2, [K<sup>29</sup>]eristicophin (4-51):

E-E-P-C-A-T-G-P-C-C-R-R-C-K-F-K-R-A-G-K-V-C-R-

V-A-K-G-D-W-N-N-D-Y-C-T-G-K-S-C-D-C-P-R-N-P-W-N-G

One-half mmol of PAM-Gly resin (0.6 meq/g, Applied Biosystems, Foster City, CA) was subjected to Procedure A with the required amino acids (introduced in order). The Boc-protected amino acids had the following side-chain protection: Arg(Tos), Asp(OcHex), Cys(4-MeBzl), Glu(OcHex), Lys(Cl-Z), Ser(OBzl), Thr(OBzl), Trp(CHO), and Tyr(Br-Z). Cleavage, refolding and purification of this peptide was identical to the previous example. Receptor

**SUBSTITUTE SHEET**

binding data for this analog are shown in Figures 26 and 28 in WO 90/15620.

#### Example 6

##### Preparation of Analog #3:

##### G-C-G-K-G-D-W-P-C-A-NH<sub>2</sub>

5 One-half mmol of pMBHA resin (0.72 meq/g, Applied Biosystems, Foster City, CA) was subjected to Procedure A with the required amino acids (introduced in order). The  
10 Boc-protected amino acids had the following side-chain protection: Asp(O-cHex), Cys(4-MeBzl), and Lys(Cl-z). Following completion of the assembly of the protected peptide-resin, the amino terminal Boc group was removed with TFA and the resin dried as its TFA-salt form. The  
15 resin (1.54 g) was treated with anhydrous hydrogen fluoride (HF) containing 10% anisole, 2% ethyl methyl sulfide for 30 min at -10°C, and an additional 30 min at 0°C. The HF was removed in vacuo and the peptide/resin mixture was suspended in diethyl ether followed by  
20 alternately washing with chloroform and ether 3X. After a final ether wash, the peptide was extracted from the resin with 2.0 M acetic acid, diluted with distilled water and lyophilized.

The crude peptide (370 mg) was dissolved in  
25 deoxygenated 10 mM NH<sub>4</sub>OAc, pH 8, to 0.5 mg/ml and allowed to oxidize by dropwise addition of a slight excess of 0.01 M potassium ferricyanide (K<sub>3</sub>Fe(CN)<sub>6</sub>) solution, stirred an additional 20 min, and adjusted to pH 5 with acetic acid. The peptide solution was treated with DOWEX  
30 AG3x4 anion-exchange resin for 15 min with stirring and the resin filtered, diluted with H<sub>2</sub>O and lyophilized to yield the crude cyclized peptide. The crude cyclized peptide (392 mg) was purified by desalting on Sephadex G-25F using 0.5M acetic acid as eluent, followed by ion-  
35 exchange chromatography on CM-Sepharose (Pharmacia) using

SUBSTITUTE SHEET

-50-

an elution gradient generated by addition of 100 mM NH<sub>4</sub>OAc to a solution of 10 mM NH<sub>4</sub>OAc, pH 4.5. Fractions which had a minimum purity of 90% by HPLC analysis were pooled and lyophilized from H<sub>2</sub>O several times to yield 175 mg. Final purification consisted of preparative HPLC purification on a Water C-18 reverse-phase column with an acetonitrile/water/TFA gradient to yield purified peptide. Receptor binding data for this analog are shown in Figures 26, 29 and 30 in WO 90/15620.

#### Example 7

##### Preparation of Additional Analogs

The following analogs were synthesized; in most cases in a manner similar to that set forth in Example 6. However, analog 60, shown below, was prepared in solution via guanidation of the side chain of the lysine residue of analog #19 using the procedure of Majusz, S., et al., FEBS Letts (1980) 110:85-87.

One mg of analog #19 was reacted with 1 mg of 1-amidino-3, 5-dimethylpyrazole nitrate (Aldrich) in 1 ml of absolute ethanol in the presence of diisopropylethylamine (DIEA) at room temperature for 4 days. The product analog 60 was purified from excess reagent and starting materials by reversed-phase HPLC on a C-18 column using a gradient of acetonitrile in 0.1% trifluoroacetic acid. Nine hundred ug of this material was isolated in purified form.

- #4 G-C-K-G-D-W-P-C-A-NH<sub>2</sub>
- #5 C-G-K-G-D-W-P-C-NH<sub>2</sub>
- #6 G-C-G-K-G-D-W-C-A-NH<sub>2</sub>
- #7 G-C-K-G-D-W-C-A-NH<sub>2</sub>
- #8 Acetyl-C-K-G-D-C-NH<sub>2</sub>
- #9 Mpr-K-G-D-Pen-NH<sub>2</sub>
- #10 C-K-G-D-W-F-C-NH<sub>2</sub>
- #11 Acetyl-C-R-G-D-Pen-NH<sub>2</sub>
- #12 C-K-G-D-Y-P-C-NH<sub>2</sub>
- #13 C-K-G-D-F-P-C-NH<sub>2</sub>
- #19 Mpr-K-G-D-W-P-C-NH<sub>2</sub>

SUBSTITUTE SHEET

-51-

- 5  
#34 C-K-G-D-W-G-C-NH<sub>2</sub>  
#35 C-K-G-E-W-P-C-NH<sub>2</sub>  
#36 C-Orn-G-D-W-P-C-NH<sub>2</sub>  
#37: C-K-A-D-W-P-C-NH<sub>2</sub>  
#38: C-K-A<sup>†</sup>-D-W-P-C-NH<sub>2</sub>  
#39: C-K-G-D-W-(Sar)-C-NH<sub>2</sub>  
#40: C-K(Formyl)-G-D-W-P-C-NH<sub>2</sub>  
#41: C-K-G-D-I-P-C-NH<sub>2</sub>  
#42: C-K-G-D-(4-Cl-Phe)-P-NH<sub>2</sub>  
10  
#43: C-K-(Sar)-D-W-P-C-NH<sub>2</sub>  
#44: C-K-G-D-(4-NO<sub>2</sub>-Phe)-P-C-NH<sub>2</sub>  
#45: C-K-G-D-(NMePhe)-P-C-NH<sub>2</sub>  
#46: C-H-G-D-W-P-C-NH<sub>2</sub>  
#47: Acetyl-C-K-G-D-W-P-C-NH<sub>2</sub>  
#48: Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>  
15  
#49: Mvl-K-G-D-W-P-C-NH<sub>2</sub>  
#50: Mpr-K-G-D-W<sup>†</sup>-P-Pen-NH<sub>2</sub>  
#51: Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>  
#52: Mpr-K-G-D-W-P-Pen<sup>†</sup>-NH<sub>2</sub>  
#53: Mpr-K-G-D-W-P<sup>†</sup>-Pen-NH<sub>2</sub>  
#54: Mpr-K-G-D<sup>†</sup>-W-P-Pen-NH<sub>2</sub>  
20  
#55: Mpr-K-G-D-W-(Thz)-C-NH<sub>2</sub>  
#56: Mpr-K-G-D-H(2,4-DNP)-P-C-NH<sub>2</sub>  
#57: Mpr-K-G-D-(2-Nal)-P-Pen-NH<sub>2</sub>  
#58: Mvl-K-G-D-W-P-Pen-NH<sub>2</sub>  
#59: Mpr-K-G-D-W-(Pip)-Pen-NH<sub>2</sub>  
25  
#60: Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>  
#61: Mpr-K-G-D-W-P-C<sup>†</sup>-NH<sub>2</sub>  
#62: Mpr-(D-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
#63: Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>  
#64: Mpr-(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
#65: Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
30  
#66: Mpr-(N<sup>G</sup>,N<sup>G'</sup>-ethylene-Har)-G-D-W-P-C-NH<sub>2</sub>  
#67: Mpr-(N<sup>G</sup>,N<sup>G'</sup>-ethylene-Har)-G-D-W-P-Pen-NH<sub>2</sub>  
#68: Mpr-Har-Sar-D-W-P-C-NH<sub>2</sub>  
#69: Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
#70: Mpr-(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
35  
#71: Mpr-Har-Sar-D-W-P-PenNH<sub>2</sub>  
#72: Mpr-(Phenylimidyl-Lys)-G-D-W-P-PenNH<sub>2</sub>  
#73: Mpr-Har-G-D-W-(3,4-dehydro P)-C-NH<sub>2</sub>

SUBSTITUTE SHEET

Example 8PAI Activity of Peptides

When tested in the standard aggregation inhibition assays described above, analogs #3-5 had IC<sub>50</sub> values of 5  $\mu$ M for ability to inhibit ADP-induced human platelet aggregation. However, analog #6 has an IC<sub>50</sub> of more than 200  $\mu$ M, and analog #7, 100  $\mu$ M. IC<sub>50</sub> values for the analogs of the invention in this assay are as follows:

	<u>Analog</u>	<u>Sequence</u>	<u>Appr. IC<sub>50</sub>(<math>\mu</math>M)</u>
	#3	G-C-G-K-G-D-W-P-C-A-NH <sub>2</sub>	5
	#4	G-C-K-G-D-W-P-C-A-NH <sub>2</sub>	5
	#5	C-G-K-G-D-W-P-C-NH <sub>2</sub>	5
	#6	G-C-G-K-G-D-W-C-A-NH <sub>2</sub>	>200
15	#7	G-C-K-G-D-W-C-A-NH <sub>2</sub>	100
	#8	Acetyl-C-K-G-D-C-NH <sub>2</sub>	200
	#9	Mpr-K-G-D-Pen-NH <sub>2</sub>	25
	#10	C-K-G-D-W-P-C-NH <sub>2</sub>	5
	#11	Acetyl-C-R-G-D-Pen-NH <sub>2</sub>	5
20	#12	C-K-G-D-Y-P-C-NH <sub>2</sub>	12
	#13	C-K-G-D-F-P-C-NH <sub>2</sub>	20
	#19	Mpr-K-G-D-W-P-C-NH <sub>2</sub>	1
	#34	C-K-G-D-W-G-C-NH <sub>2</sub>	100
	#35	C-K-G-E-W-P-C-NH <sub>2</sub>	>300
	#36	C-Orn-G-D-W-P-C-NH <sub>2</sub>	150-200
25	#37	C-K-A-D-W-P-C-NH <sub>2</sub>	100
	#38	C-K-A <sup>†</sup> -D-W-P-C-NH <sub>2</sub>	>200
	#39	C-K-G-D-W-(Sar)-C-NH <sub>2</sub>	5
	#40	C-K(Formyl)-G-D-W-P-C-NH <sub>2</sub>	>200
	#41	C-K-G-D-I-P-C-NH <sub>2</sub>	100
30	#42	C-K-G-D-(4-Cl-Phe)-P-NH <sub>2</sub>	20
	#43	C-K-(Sar)-D-W-P-C-NH <sub>2</sub>	50
	#44	C-K-G-D-(4-NO <sub>2</sub> -Phe)-P-C-NH <sub>2</sub>	75
	#45	C-K-G-D-(NMePhe)-P-C-NH <sub>2</sub>	>200
	#46	C-H-G-D-W-P-C-NH <sub>2</sub>	200

-53-

	#47	Acetyl-C-K-G-D-W-P-C-NH <sub>2</sub>	2.5
	#48	Mpr-K-G-D-W(Formyl)-P-C-NH <sub>2</sub>	1
	#49	Mvl-K-G-D-W-P-C-NH <sub>2</sub>	1.5
5	#50	Mpr-K-G-D-W <sup>†</sup> -P-Pen-NH <sub>2</sub>	>200
	#51	Mpr-K-G-D-W-P-Pen-NH <sub>2</sub>	0.75
	#52	Mpr-K-G-D-W-P-Pen <sup>†</sup> -NH <sub>2</sub>	5
	#53	Mpr-K-G-D-W-P <sup>†</sup> -Pen-NH <sub>2</sub>	>200
	#54	Mpr-K-G-D <sup>†</sup> -W-P-Pen-NH <sub>2</sub>	>100
	#55	Mpr-K-G-D-W-(Thz)-C-NH <sub>2</sub>	2
10	#56	Mpr-K-G-D-H(2,4-DNP)-P-C-NH <sub>2</sub>	5
	#57	Mpr-K-G-D-(2-Nal)-P-Pen-NH <sub>2</sub>	1
	#58	Mvl-K-G-D-W-P-Pen-NH <sub>2</sub>	1
	#59	Mpr-K-G-D-W-(Pip)-Pen-NH <sub>2</sub>	1
	#60	Mpr-(Har)-G-D-W-P-C-NH <sub>2</sub>	0.15
15	#61	Mpr-K-G-D-W-P-C <sup>†</sup> -NH <sub>2</sub>	15
	#62	Mpr-K <sup>†</sup> -G-D-W-P-Pen-NH <sub>2</sub>	2.5
	#63	Mpr-(Har)-G-D-W-P-Pen-NH <sub>2</sub>	0.10
	#64	Mpr-(Acetimidyl-Lys)-G-D-W-P-C-NH <sub>2</sub>	0.25
	#68	Mpr-Har-Sar-D-W-P-C-NH <sub>2</sub>	3.0
	#69	Mpr-(Acetimidyl-Lys)-G-D-W-P-Pen-NH <sub>2</sub>	0.5
20	#70	Mpr-(Phenylimidyl-Lys)-G-D-W-P-C-NH <sub>2</sub>	0.5
	#71:	Mpr-Har-Sar-D-W-P-PenNH <sub>2</sub>	2.5
	#72:	Mpr-(Phenylimidyl-Lys)-G-D-W-P-PenNH <sub>2</sub>	0.5

25

Example 9Activity of Linear versus Cyclic Peptides

When tested for inhibition of fibrinogen binding to  
 30 GP IIb-IIIa in the plate assay, linear RGDW-NH<sub>2</sub> was very  
 similar in activity to cyclic GCGRGDWPCA-NH<sub>2</sub>. In  
 contrast, the linear KGDW-NH<sub>2</sub> was much less potent than  
 cyclic GCGKGDWPCA-NH<sub>2</sub>. For the KGDW compounds, but not  
 the RGDW compounds, cyclization resulted in a marked

35

SUBSTITUTE SHEET

-54-

increase in the ability of the peptide to inhibit the binding of fibrinogen to GP IIb-IIIa.

#### Example 10

5

##### Results of Plate Binding

##### Assays for Synthetic Peptides

The peptides synthesized in Example 8, in addition to being assessed for the ability to inhibit platelet aggregation directly, were also tested in the plate assays as described above and indicated that these analogs are differentially capable, to varying degrees, of inhibiting the binding of fibrinogen to GP IIb-IIIa as compared to vitronectin to vitronectin receptor. Analog #4 appears, among this group, to have the highest differential. Analogs #7 and #5, on the other hand, are also quite specific, and have excellent platelet aggregation inhibition activities.

#### Example 11

20

##### Effects of Purified Peptides on Cell Adhesion

M21 melanoma cells were labelled with <sup>35</sup>S-methionine, and then added to vitronectin-coated plates in the presence of the indicated concentrations of purified snake venom peptides. Cell attachment was measured by solubilizing the cells remaining after an incubation and wash, as previously describe. Neither barbourin nor Peptide 1 (truncated barbourin) had a significant effect on cell adhesion to vitronectin, although both are potent inhibitors of platelet aggregation. In contrast, cotiarin, which is a potent inhibitor of vitronectin binding to the vitronectin receptor, was very potent in inhibiting cell attachment to vitronectin. In similar experiment, Peptide #3, Peptide #3 with K replaced by R (GCGRGDWPCA-NH<sub>2</sub>) and RGDS were examined on M21 cell attachment to vitronectin.

SUBSTITUTE SHEET



-55-

RGDS and GCGRGDWPCA-NH<sub>2</sub> are potent inhibitors of cell attachment whereas GCGKGDWPCA-NH<sub>2</sub> was ineffective up to 60 uM.

5

Example 12Comparison of Analogs 60 and 19

Analogs 60 and 19 described above containing the sequence K<sup>\*</sup>GDX and are identical except for the embodiment of K<sup>\*</sup>. Analog 60 is of the formula:

10

Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>;

analog 19 is of the formula:

Mpr-K-G-D-W-P-C-NH<sub>2</sub>.

15

These analogs were tested by standard platelet aggregation inhibition assays and using the cell adhesion assay of Example 11 above. Analog #60 was efficient at vanishingly small concentrations in inhibiting platelet aggregation, and was relatively less effective in preventing cell adhesion to vitronectin. Analog #19 had good platelet aggregation inhibition activity as well as specificity; however, it was less active in the platelet aggregation inhibition assay than its analog #60 counterpart. Analog #60 had an IC<sub>50</sub> in platelet aggregation of approximately 0.15 nM; analog #19 had an IC<sub>50</sub> of approximately 1 nM.

20

25

Example 1330 Construction of Expression Vectors for Barbourin Peptides

A gene encoding the full length [L<sup>41</sup>] barbourin peptide (1-73) was assembled from synthetic oligonucleotides as shown in figure 38 of WO 90/15620, which were kinased, annealed and ligated into EcoRI-HindIII digested M13mp18 using standard procedures. The

35

SUBSTITUTE SHEET

-56-

bacterial alkaline phosphatase gene (phoA) signal sequence (Watson, M.E.E., Nucleic Acids Research (1984) 12:5145) was added to the barbourin construct by ligating synthetic oligonucleotides into the EcoRI/NcoI sites of the [L<sup>41</sup>] barbourin (1-73) construct as shown in Figure 39 of WO 90/15620. The nucleotide sequences of all constructs were verified by the Sanger dideoxy chain termination method.

A truncated version of this peptide was also constructed from synthetic oligonucleotides which would encode only amino acids 28-73 of the full length molecule. Two alterations, Q<sup>28</sup> to E<sup>28</sup> and A<sup>64</sup> to C<sup>64</sup> were introduced using site directed mutagenesis as described by Kunkel et al. Meth Enzymol (1987) 154:367. The phoA signal sequence was added to the truncated version as described above (Figure 40 of WO 90/15620). In addition, the signal sequence for the E. coli heat-stable enterotoxin II (Picken, R.W., et al. Infect Immun (1983) 42:269) was added to the truncated version using synthetic oligonucleotides with EcoRI and NcoI compatible ends. All bacterial secretion constructs were subcloned into the bacterial expression vector pPROK-1 (Brosius, J., Gene (1984) 27:151, *ibid*:161), available commercially from CLONTECH Lab, Inc. using EcoRI and HindIII restriction endonucleases.

A gene encoding tandem repeats of the desired title peptide was prepared using the polymerase chain reaction (PCR) to produce the multimerization unit from the full-length barbourin peptide 1-73 containing L41 and C64.

Figure 41 of WO 90/15620 shows the oligonucleotides used for the PCR synthesis. The PCR reaction was conducted according to the method of Saki, R.K., et al. Science (1988) 239:487. The resulting polymer junction contains methionines at either end of the sequence as

35

SUBSTITUTE SHEET

-57-

shown in WO 90/15620, Figure 42, and provides desirable restriction sites for the construct.

The tandem repeats are formed from the individual multimer-forming components by, for example, ligating an  
5 EcoRI/BamHI fragment to a BglII/HindIII fragment in an M13mp18 vector cut with EcoRI/HindIII to form a dimer. The resultant dimer is excised with EcoRI and BamHI and relegated to a BglII/HindIII fragment to produce a trimer, and so on until the desired size is obtained.  
10 This construction is diagramed in Figure 43 of WO 90/15620.

The multimer was then ligated into the E. coli vector pKK233-2, Amann, E., et al., Gene (1985) 40:183, available from Clontech, by digesting the vector with  
15 NcoI/HindIII and ligating a monomer subfragment of NcoI/BamHI and multimer subfragments of BglII/HindIII.

For expression as a fusion protein, the above digested vector was used along with an NcoI/EcoRI subfragment containing a slightly modified amino-terminal  
20 portion(amino acids 1 to 72) of the chloramphenicol acetyltransferase gene (Chang, C.N., et al., Gene (1987) 55:189) and EcoRI-HindIII subfragments of the multimer constructions.

25

#### Example 14

##### Expression of Recombinant Genes

Protein expression from all of the recombinant plasmids described above is induced according to Kanamari  
et al., Gene (1988) 66:295, after transfection into  
30 appropriate E. coli host strains. Products are characterized by sodium dodecyl sulfate polyacrylamide gel electrophoresis and by their ability to inhibit ADP-induced platelet aggregation in platelet-rich plasma. Following purification, the multimeric proteins are

35

SUBSTITUTE SHEET

-58-

converted to monomer units with cyanogen bromide cleavage and the products assayed as above.

Example 155                    Preparation of Additional Analogs

The following analogs were synthesized in a manner similar to that set forth in Example 6 and were tested for PAI activity in the assay method described hereinabove.

10			<u>Appr. IC<sub>50</sub></u> <u>(<math>\mu</math>M)</u>
	<u>Analog</u>	<u>Sequence</u>	
	PAI 80	Mpr-P-Har-G-D-W-P-C-NH <sub>2</sub>	1.29
	PAI 81	Mpr-G-Har-G-D-W-P-C-NH <sub>2</sub>	7.47
	PAI 82	Mpr-A-Har-G-D-W-P-C-NH <sub>2</sub>	0.12
15	PAI 83	Mpr-Aib-Har-G-D-W-P-C-NH <sub>2</sub>	2.20
	PAI 84	Mpr-(N-Me-Arg)-Har-G-D-W-P-C-NH <sub>2</sub>	0.25
	PAI 85	Mpr-(N-Me-Ser)-Har-G-D-W-P-C-NH <sub>2</sub>	0.28
	PAI 86	Mpr-A <sup>†</sup> -Har-G-D-W-P-C-NH <sub>2</sub>	1.15
	PAI 87	Mpr-( $\beta$ -Ala)-Har-G-D-W-P-C-NH <sub>2</sub>	0.92
20	PAI 88	Mpr-(N-Me-Leu)-Har-G-D-W-P-C-NH <sub>2</sub>	0.84
	PAI 89	Mpr-(N-Me-Ala)-Har-G-D-W-P-C-NH <sub>2</sub>	0.62
	PAI 90	Mpr-Sar-Har-G-D-W-P-C-NH <sub>2</sub>	0.27
	PAI 91	Mpr-V-Har-G-D-W-P-C-NH <sub>2</sub>	0.35
	PAI 92	Mpr-S-Har-G-D-W-P-C-NH <sub>2</sub>	0.24
25	PAI 93	Mpr-Har-G-D-W-P-A-C-NH <sub>2</sub>	3.33
	PAI 94	Mpr-Har-G-D-W-P-(N-Me-Ala)-C-NH <sub>2</sub>	1.46
	PAI 95	Mpr-Har-G-D-W-P-G-C-NH <sub>2</sub>	8.66
	PAI 96	Mpr-Har-G-D-W-P-A <sup>†</sup> -C-NH <sub>2</sub>	0.23
	PAI 97	Mpr-Har-G-D-W-P-P-C-NH <sub>2</sub>	1.40
30	PAI 98	Mpr-Har-G-D-W-P-(Sar)-C-NH <sub>2</sub>	0.31
	PAI 99	Mpr-Har-G-D-W-P-(Aib)-NH <sub>2</sub>	0.46
	PAI 100	Mpr-A-(Har)-G-D-W-P-Pen-NH <sub>2</sub>	0.37
	PAI 101	Mpr-A-K-G-D-W-P-Pen-NH <sub>2</sub>	4.91
	PAI 102	Mpr-D-(Har)-G-D-W-P-Pen-NH <sub>2</sub>	4.04

35

Example 16 Stability Determination

A solution was formed in 1.0 M citric acid of greater than 95% pure cyclic form of Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate polypeptide (I) at a concentration of up to 200 mg per ml of solution. This solution was diluted to the desired final peptide concentration as follows: This solution was first diluted to about 85% of the final volume with water. The pH of the solution was adjusted to 5.0-5.5 using sodium hydroxide. The solution was then diluted to the final volume with water. Compositions having a concentration of (I) of 0.5 mg/ml and 5.0 mg/ml were formulated and evaluated for stability by HPLC, UV and visual inspection as described in Example 1. The compositions were also evaluated for platelet aggregation inhibiting activity using the assay described above. Results of these experiments are set forth in Tables 4 and 5 below.

Table 4

Stability of 0.5 mg/ml Composition

	Temp (°C)	Time (Days)	CAC <sup>a</sup>	pH	HPLC <sup>b</sup> (%Area)	HPLC (mg/mL)	UV (mg/mL)	Activity <sup>c</sup> (nM)
25	5	0	CC	5.44	98.6	0.51	0.50	175
	5	32	CC	5.45	98.8	0.50	0.49	183
	-15	32	CC	5.46	98.9	0.50	0.50	199
	30	32	CC	5.44	98.4	0.50	0.50	174
	45	32	CC	5.44	97.1	0.50	0.50	202

<sup>a</sup> CC=Clear, colorless solution; CvPY=Clear, very pale yellow solution;

<sup>b</sup> Purity expressed as a percent of total peak area.

<sup>c</sup> Activity expressed as an IC<sub>50</sub> (the concentration calculated to achieve a 50% inhibition of platelet aggregation).

SUBSTITUTE SHEET

-60-

Table 5

Stability of 5 mg/ml Composition

	Temp (°C)	Time (Days)	CAC <sup>a</sup>	pH	HPLC <sup>b</sup> (%Area)	HPLC (mg/mL)	UV (mg/mL)	Activity <sup>c</sup> (nM)
5	5	0	CC	5.39	99.5	5.30	5.04	88
	5	29	CC	5.40	98.3	5.35	NT	NT
10	5	60	CC	5.40	98.3	4.71	NT	NT
	5	90	CC	5.38	97.9	4.97	NT	NT
	30	29	CC	5.41	98.4	4.76	NT	NT
	30	60	CC	5.40	97.4	4.64	NT	NT
15	30	90	CC	5.39	96.9	4.87	NT	NT
	45	29	CC	5.41	96.4	4.65	NT	NT
	45	69	CC	5.41	93.9	4.22	NT	NT
	45	90	CvPY	5.39	89.8	4.14	NT	NT

a CC=Clear, colorless solution; CvPY=Clear, very pale yellow solution;

b Purity expressed as a percent of total peak area.

c Activity expressed as an IC<sub>50</sub> (the concentration calculated to achieve a 50% inhibition of platelet aggregation).

NT No testing required at this timepoint.

Results of these experiments set forth in Tables 4 and 5 demonstrated that the liquid solutions of the polypeptide of the invention having a pH of 5.0-5.5 were very stable from about -15°C to about 45°C. The compositions were also found to be normal on visual inspection.


SUBSTITUTE SHEET

-61-


What is claimed is:

1. A method of storage-stabilizing a substantially pure polypeptide by dissolving said polypeptide in a citrate buffer to form a storage-stable solution having a pH of from about 5.0 to about 5.5.

2. A method of storage-stabilizing a substantially pure polypeptide comprises forming a liquid solution consisting essentially of a substantially pure polypeptide by dissolving said polypeptide in a citrate buffer to storage-stabilize said polypeptide in said solution, said solution having a pH of from about 5.0 to about 5.5, said polypeptide has the formula



wherein  $K^*$  has the formula



wherein (G/Sar) is G or Sar;

wherein each  $R^1$  is independently H, alkyl(1-6C), or at most one  $R^1$  is  $R^2-C=NR^3$ ,


wherein  $R^2$  is H, alkyl(1-6C) or is a substituted or unsubstituted phenyl or benzyl residue, or is  $NR^4_2$  in which each  $R^4$  is independently H or alkyl (1-6C), and  $R^3$  is H, alkyl (1-6C), phenyl or benzyl, or  $R^2-C=NR^3$  is a radical selected from the group

consisting of

35

SUBSTITUTE SHEET

-62-



5

where m is an integer of 2-3, and each R<sup>5</sup> is  
 10 independently H or alkyl(1-6C);

and where one or two (CH<sub>2</sub>) may be replaced by O or S  
 provided said O or S is not adjacent to another  
 heteroatom;

AA<sub>1</sub> is a small, neutral (polar or nonpolar) amino  
 15 acid and n1 is an integer of 0-3;

AA<sub>2</sub> is a neutral, nonpolar large (aromatic or  
 nonaromatic) or a polar aromatic amino acid and n2 is an  
 integer of 0-3);

AA<sub>3</sub> is a proline residue or a modified proline  
 20 residue and n3 is an integer of 0-1;

AA<sub>4</sub> is a neutral, small amino acid or the  
 N-alkylated form thereof and n4 is an integer of 0-3;

each of X<sub>1</sub> and X<sub>2</sub> is independently a residue capable  
 of forming a bond between X<sub>1</sub> and X<sub>2</sub> to obtain a cyclic  
 25 compound as shown; and

each of Y<sub>1</sub> and Y<sub>2</sub> is independently a noninterfering  
 substituent or may be absent;

wherein one or more peptide linkages may optionally  
 be replaced by linkage selected from the group consisting  
 30 of -CH<sub>2</sub>NH-, -CH<sub>2</sub>S-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- (cis and trans),  
 -COCH<sub>2</sub>-, -CH(OH)CH<sub>2</sub>- and -CH<sub>2</sub>SO-.

3. The method of claim 2 wherein said polypeptide  
 Y<sub>1</sub> is H, acyl, or a peptide residue or derivatized form  
 35 thereof or is absent and Y<sub>2</sub> is OH, NH<sub>2</sub> or a peptide



-63-

residue or derivatized form thereof or is absent,  $Y_2$  is  $NH_2-A-NH_2$  or is absent,  $X_1$  and  $X_2$  are selected from the group consisting of cysteine (C), mercaptopropionyl (Mpr) and penicillamine (Pen),  $AA_1$  is G and  $n_1$  is 0 or 1,  $AA_2$  is selected from the group consisting of W, F, L, Y, and V and  $K^*$  is K, Har, acetimidyl-Lys or phenylimidyl-Lys.

4. The method of claim 3 wherein the polypeptide is selected from

10 Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>  
 Mvl-K-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>  
 15 Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 20 Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-K-G-D-W-P-C-NH<sub>2</sub>  
 or cyclic forms thereof.

5. The method of claim 2 wherein the polypeptide is  
 25 Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub> or a cyclic form thereof.

6. The method of claim 2 wherein the polypeptide is  
 Mvl-K-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

30 7. The method of claim 2 wherein the polypeptide is  
 Mpr-K-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

8. The method of claim 2 wherein the polypeptide is  
 Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

35

-64-

9. The method of claim 2 wherein the polypeptide is Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

10. The method of claim 2 wherein the polypeptide is Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

11. The method of claim 2 wherein the polypeptide is Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

12. The method of claim 2 wherein the polypeptide is Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

13. The method of claim 2 wherein the polypeptide is Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

14. The method of claim 2 wherein the polypeptide is Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

15. The method of claim 2 wherein the polypeptide is Mpr-K-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

16. A method according to claim 2 wherein said storage stability at about 70°C or less for at least 49 days.

17. A method of storage-stabilizing an injectable biologically active substantially pure polypeptide comprising forming a liquid solution consisting essentially of a biologically active substantially pure polypeptide by dissolving said polypeptide in a citrate buffer, said solution having a pH of from about 5.0 to

SUBSTITUTE SHEET


-65-

about 5.5, to storage-stabilize said polypeptide in said solution.

18. The method according to claim 17 wherein said  
5 polypeptide is a synthetic or recombinant polypeptide.


19. The method according to claim 18 wherein said  
polypeptide is biologically active for inhibiting  
thrombus formation, preventing platelet loss during  
10 extracorporeal circulation of blood or for treating a  
patient suspected of having a platelet-associated  
ischemic syndrome.

20. The method according to claim 3 wherein said  
15 polypeptide has the formula



wherein  $K^*$  has the formula

20




wherein (G/Sar) is G or Sar;


wherein each  $R^1$  is independently H, alkyl(1-6C), or  
25 at most one  $R^1$  is  $R^2-C=NR^3$ ,

wherein  $R^2$  is H, alkyl(1-6C) or is a substituted or  
unsubstituted phenyl or benzyl residue, or is  $NR^4_2$  in  
which each  $R^4$  is independently H or alkyl(1-6C), and

$R^3$  is H, alkyl(1-6C), phenyl or benzyl, or  
30  $R^2-C=NR^3$  is a radical selected from the group  
consisting of



. and



35

SUBSTITUTE SHEET

-66-

where m is an integer of 2-3, and each R<sup>5</sup> is independently H or alkyl(1-6C);

and wherein one or two (CH<sub>2</sub>) may be replaced by O or S provided said O or S is not adjacent to another heteroatom;

AA<sub>1</sub> is a small, neutral (polar or nonpolar) amino acid and n<sub>1</sub> is an integer of 0-3;

AA<sub>2</sub> is a neutral, nonpolar large (aromatic or nonaromatic) or a polar aromatic amino acid and n<sub>2</sub> is an integer of 0-3;

AA<sub>3</sub> is a proline residue or a modified proline residue and n<sub>3</sub> is an integer of 0-1;

AA<sub>4</sub> is a neutral, small amino acid or the N-alkylated form thereof and n<sub>4</sub> is an integer of 0-3;

each of X<sub>1</sub> and X<sub>2</sub> is independently a residue capable of forming a bond between X<sub>1</sub> and X<sub>2</sub> to obtain a cyclic compound as shown; and

each of Y<sub>1</sub> and Y<sub>2</sub> is independently a noninterfering substituent or may be absent;

wherein one or more peptide linkages may optionally be replaced by a linkage selected from the group consisting of -CH<sub>2</sub>NH-, -CH<sub>2</sub>S-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH- (cis and trans), -COCH<sub>2</sub>-, -CH(OH)CH<sub>2</sub>- and -CH<sub>2</sub>SO-.

21. The method according to claim 19 wherein the polypeptide is a cyclic polypeptide containing up to 10 amino acids residues.

22. The method according to claim 21 wherein the polypeptide is Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate or a cyclic form thereof.

23. The method according to claim 17 wherein the pH is about 5.25.

SUBSTITUTE SHEET

-67-


24. The method according to claim 23 wherein the stabilizing is at about -15°C to about 30°C.

25. The method according to claim 22 wherein the stability is at about 5 to about 30°C.

26. The method according to claim 17 wherein said polypeptide is a cyclic polypeptide containing up to 10 amino acid residues and is stabilized in a solution having a pH of about 5.25, in which the citrate buffer solution includes sodium hydroxide.


27. A storage-stable composition consisting essentially of a substantially pure polypeptide dissolved in a citrate buffer, said solution having a pH of from about 5.0 to about 5.5.

28. A storage-stable composition consisting essentially of a effective amount of a substantially pure polypeptide dissolved in a citrate buffer, said solution having a pH of from about 5.0 to about 5.5., wherein the polypeptide has the formula



25

wherein  $K^*$  has the formula



30

wherein (G/Sar) is G or Sar;

wherein each  $R^1$  is independently H, alkyl(1-6C), or at most one  $R^1$  is  $R^2-C=NR^3$ ,


35

wherein  $R^2$  is H, alkyl(1-6C) or is a substituted or unsubstituted phenyl or benzyl residue, or is  $NR^4_2$  in which each  $R^4$  is independently H or alkyl(1-6C), and

SUBSTITUTE SHEET

-68-

$R^3$  is H, alkyl(1-6C), phenyl or benzyl, or  
 $R^2-C=NR^3$  is a radical selected from the group  
 consisting of



10 where m is an integer of 2-3, and each  $R^5$  is  
 independently H or alkyl (1-6C);

and wherein one or two  $(CH_2)$  may be replaced by O or  
 S provided said O or S is not adjacent to another  
 heteroatom;

15  $AA_1$  is a small, neutral (polar or nonpolar) amino  
 acid and  $n_1$  is an integer of 0-3;

$AA_2$  is a neutral, nonpolar large (aromatic or  
 nonaromatic) or a polar aromatic amino acid and  $n_2$  is an  
 integer of 0-3;

20  $AA_3$  is a proline residue or a modified proline  
 residue and  $n_3$  is an integer of 0-1;

$AA_4$  is a neutral, small amino acid of the N-  
 alkylated form thereof and  $n_4$  is an integer of 0-3;

25 each of  $X_1$  and  $X_2$  is independently a residue capable  
 of forming a bond between  $X_1$  and  $X_2$  to obtain a cyclic  
 compound as shown; and

each of  $Y_1$  and  $Y_2$  is independently a noninterfering  
 substituent or may be absent;

30 wherein one or more peptide linkages may optionally  
 be replaced by a linkage selected from the group  
 consisting of  $-CH_2NH-$ ,  $-CH_2S-$ ,  $-CH_2CH_2-$ ,  $-CH=CH-$  (cis and  
 trans),  $-COCH_2-$ ,  $-CH(OH)CH_2-$  and  $-CH_2SO-$ .

29. The composition of claim 28 wherein said  
 polypeptide  $Y_1$  is H, acyl, or a peptide residue or  
 35 derivatized form thereof or is absent and  $Y_2$  is OH,  $NH_2$

-69-

or a peptide residue or derivatized form thereof or is absent,  $Y_2$  is  $NH_2-A-NH_2$  or is absent,  $X_1$  and  $X_2$  are selected from the group consisting of cysteine (C), mercaptopropionyl (Mpr) and penicillamine (Pen),  $AA_1$  is G and  $n_1$  is 0 or 1,  $AA_2$  is selected from the group consisting of W, F, L, Y, and V and  $K^*$  is K, Har, acetimidyl-Lys or phenylimidyl-Lys.

30. The composition of claim 28 wherein the polypeptide is selected from

Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub>  
 Mvl-K-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-K-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr(Phenylimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub>  
 Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub>  
 Mpr-K-G-D-W-P-C-NH<sub>2</sub>  
 or cyclic forms thereof.

31. The composition of claim 28 wherein the polypeptide is Mpr-K-G-D-W(Formyl)-P-C-NH<sub>2</sub> or a cyclic form thereof.

32. The composition of claim 28 wherein the polypeptide is Mvl-K-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

33. The composition of claim 28 wherein the polypeptide is Mpr-K-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

SUBSTITUTE SHEET

-70-

34. The composition of claim 28 wherein the polypeptide is Mpr-(Har)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

5        35. The composition of claim 28 wherein the polypeptide is Mpr-(Har)-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

10       36. The composition of claim 28 wherein the polypeptide is Mpr(Acetimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

15       37. The composition of claim 28 wherein the polypeptide is Mpr(Acetimidyl-Lys)-G-D-W-P-Pen-NH<sub>2</sub> or a cyclic form thereof.

20       38. The composition of claim 28 wherein the polypeptide is Mpr(Phenylimidyl-Lys)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

39. The composition of claim 28 wherein the polypeptide is Mpr-Ala-(Har)-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

25       40. The composition of claim 28 wherein the polypeptide is Mpr-L-homoarginine-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

30       41. The composition of claim 28 wherein the polypeptide is Mpr-K-G-D-W-P-C-NH<sub>2</sub> or a cyclic form thereof.

42. A composition of claim 28 stabilized at about 70°C or less for at least 49 days.

35

SUBSTITUTE SHEET




-71-

43. A storage-stable therapeutic liquid composition consisting essentially of an injectable biologically effective amount of a substantially pure polypeptide dissolved in a liquid solution of a citrate buffer, said  
5 solution having a pH of from about 5.0 to about 5.5.


44. The liquid composition according to claim 43 wherein said polypeptide is a synthetic or recombinant polypeptide.  
10

45. The liquid composition according to claim 44 wherein said polypeptide is biologically active for inhibiting thrombus formation, preventing platelet loss during extracorporeal circulation of blood or for  
15 treating a patient suspected of having a platelet-associated ischemic syndrome.

46. The liquid composition according to claim 45 wherein the polypeptide has the formula  
20



wherein  $K^*$  has the formula



wherein (G/Sar) is G or Sar;

wherein each  $R^1$  is independently H, alkyl(1-6C), or at most one  $R^1$  is  $R^2-C=NR^3$ ,


30 wherein  $R^2$  is H, alkyl(1-6C) or is a substituted or unsubstituted phenyl or benzyl residue, or is  $NR^4_2$  in which each  $R^4$  is independently H or alkyl(1-6C), and  $R^3$  is H, alkyl(1-6C), phenyl or benzyl, or

35

SUBSTITUTE SHEET

-72-

$R^2-C=NR^3$  is a radical selected from the group consisting of



where  $m$  is an integer of 2-3, and each  $R^5$  is independently H or alkyl (1-6C);

10 and wherein one or two  $(CH_2)$  may be replaced by O or S provided said O or S is not adjacent to another heteroatom;

$AA_1$  is a small, neutral (polar or nonpolar) amino acid and  $n_1$  is an integer of 0-3;

15  $AA_2$  is a neutral, nonpolar large (aromatic or nonaromatic) or a polar aromatic amino acid and  $n_2$  is an integer of 0-3;

$AA_3$  is a proline residue or a modified proline residue and  $n_3$  is an integer of 0-1;

20  $AA_4$  is a neutral, small amino acid or the N-alkylated form thereof and  $n_4$  is an integer of 0-3;

each of  $X_1$  and  $X_2$  is independently a residue capable of forming a bond between  $X_1$  and  $X_2$  to obtain a cyclic compound as shown; and

25 each of  $Y_1$  and  $Y_2$  is independently a noninterfering substituent or may be absent;

wherein one or more peptide linkages may optionally be replaced by a linkage selected from the group consisting of  $-CH_2NH-$ ,  $-CH_2S-$ ,  $-CH_2CH_2-$ ,  $-CH=CH-$  (cis and trans),  $-COCH_2-$ ,  $-CH(OH)CH_2-$  and  $-CH_2SO-$ .

35 47. The liquid composition according to claim 44 wherein said polypeptide is a cyclic polypeptide of up to 10 amino acids containing up to 10 amino acid residues.

-73-

48. The liquid composition according to claim 47 wherein the polypeptide is Mpr-L-homoarginine-Gly-Asp-Trp-Pro-Cys-NH<sub>2</sub> • acetate or a cyclic form thereof.

5 49. The composition according to claim 43 wherein said pH is about 5.25.

10 50. The liquid composition according to claim 43 wherein said polypeptide is a cyclic polypeptide containing up to 10 amino acid residues and is stabilized in a solution having a pH of about 5.25, in which the citrate buffer solution includes sodium hydroxide.

15 51. A composition comprising a substantially pure polypeptide and a stabilizing effective amount of a citrate buffer, said composition having a pH of from about 5.0 to about 5.5.

20 52. An article of manufacture comprising a sterile delivery vial, bag or bottle filled with a liquid composition of claim 43 in injectable form.

25

30

35

SUBSTITUTE SHEET

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/03933

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07K1/00; C07K7/06; A61K37/02		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	C07K ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	WO,A,8 304 027 (AMERICAN NATIONAL RED CROSS) 24 November 1983 * summary of the invention, pages 3-4 *	1-52
Y	METH. ENZYMOL. vol. I, 1955, pages 138 - 146 GOMORI, G. (COLOWICK & KAPLAN EDS.) ACADEMIC PRESS 'Preparation of buffers for use in enzyme studies' * pages 140 and 141 *	1-52
<p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
09 JULY 1993		12.08.93
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		HERMANN R.

Form PCT/ISA/210 (second sheet) (January 1985)

## III. DOCUMENTS CONSIDERED TO BE RELEVANT

(CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	METH. ENZYMOL. vol. I, 1955, pages 67 - 79 ALDEN-GREEN, A. & HUGHES, W. L. 'Protein fractionation on the basis of solubility in aqueous solutions ...' * page 75 *	1-52
Y	----- LEHNINGER, A.L. 'Biochemistry' 1975 , WORTH PUBLISHERS, INC. * page 162 * -----	1-52

US 9303933  
SA 74071

09/07/93

**EPQ FORM P0479**

\\SDOCID: <WO\_\_\_\_\_.9322335A1\_I\_>